

**MARSH PLANT ASSOCIATIONS OF
SOUTH SAN FRANCISCO BAY:
1999 COMPARATIVE STUDY**

Prepared by

H. T. HARVEY & ASSOCIATES

Ronald R. Duke, M.A., Principal
Patrick J. Boursier, Ph.D., Senior Plant Ecologist
Eric Webb, Ph.D., Project Manager
Max Busnardo, M.S., Restoration Ecologist
John Bourgeois, M. S. Wetlands Ecologist
Andrew Dilworth, B. S., Plant Ecologist

Prepared for

City of San Jose
Environmental Services Department
San Jose/Santa Clara Water Pollution Control Plant
700 Los Esteros Road
San Jose, CA 95134

November 29, 1999

Project No. 477-22

TABLE OF CONTENTS

TABLE OF CONTENTS	i
EXECUTIVE SUMMARY	1
INTRODUCTION	3
SURVEY METHODS	8
STUDY AREA	8
AERIAL PHOTOGRAPHY AND ORTHORECTIFICATION	8
VEGETATION ASSOCIATION MAPPING AND AREA CALCULATIONS	9
VEGETATION ASSOCIATION CATEGORIZATION METHODS	11
AREA COMPARISONS	15
RESULTS	16
GENERAL SPECIES DISTRIBUTION, DOMINANT SPECIES CATEGORY AND HABITAT ACREAGES FOR 1999	16
Main Study Area	16
Reference Area (Alviso Slough)	20
TEMPORAL AND SPATIAL CHANGES IN MARSH HABITAT ACREAGES FROM 1989 THROUGH 1999	21
New Marsh Formation (Salt, Brackish, and Freshwater Marsh Combined)	21
Changes in Surface Area of Salt, Brackish, and Freshwater Marsh Habitats	21
Temporal Changes in Proportional Area of Salt and Brackish Marsh Between the Main Study and Reference Areas	27
Habitat Type Conversion	30
DISCUSSION	31
REFERENCES	33
APPENDIX A SOUTH BAY MARSHES: 1999 VEGETATION MAPS	36
APPENDIX B VEGETATION MATRICES	43
APPENDIX C SOUTH BAY MARSHES: PLANT LIST	62

FIGURES:

Figure 1. Segment Locations	6
Figure 2. Location of Four Reaches	7
Figure 3. Dominant Species Categories of the Main Study Area by Habitat Type, 1999	18
Figure 5. Total Marsh Acreage Comparison From 1989 to 1999 by Reach	25
Figure 6. Salt Marsh Acreage Comparison From 1989 to 1999 by Reach	26
Figure 7. Brackish Marsh Acreage Comparison From 1989 to 1999 by Reach	26
Figure 8. Freshwater Marsh Acreage Comparison From 1989 to 1999 by Reach	27
Figure 9. Temporal Comparison of the Proportion of Salt Marsh Area Between the Main Study and Reference Areas	28
Figure 10. Temporal Comparison of the Proportion of Brackish Marsh Area Between the Main Study and Reference Areas	28

TABLES:

[Table 1. Segment Names and Numbers.....](#)10

[Table 2. Dominant Species Categories, Marsh Type, and Vegetation Associations for 1989, 1998 and 1999.....](#)12

[Table 3. Summary of Acreages of the Main Study Area by dominant species categories for each habitat type for 1999.....](#)17

[Table 4. Summary of Acreages of the Reference Area \(Alviso Slough\) by dominant species categories for each habitat type for 1998 and 1999.](#)20

[Table 5. Summary of Acreages of the Main Study Area by dominant species categories for each habitat type for 1989 and 1999.](#)23

[Table 6. Summary of Acreages of the Reference Area \(Alviso Slough\) by dominant species categories for each habitat type for 1989 and 1999.](#)24

[Table 7. Detailed Evaluation of Changes in Acreage for Segment Locations by Habitat Type and for Total Marsh, 1989 to 1999.....](#)29

EXECUTIVE SUMMARY

This study is a continuation of a long-term monitoring program designed to detect changes in habitat types within the coastal marshes of South San Francisco Bay. Additionally, this study evaluates the possible contribution of the freshwater discharge from the San Jose/Santa Clara WPCP on the distribution of these habitat types. To determine the distribution of habitat types in the South Bay Marshes, the study area was subdivided into three zones: the upper segments, the transition segments and the lower segments. To track the changes that could potentially be caused by anthropogenic influences instead of environmental influences, a Reference Area (Alviso Slough) was also mapped. Vegetation mapping within the Main Study and Reference Areas commenced in 1989 and subsequent mapping has occurred in 1991, 1994, 1995, 1996, 1997 and 1998.

The 1999 plant association mapping was done on digital orthos images created from rectified color infrared aerial photography. All vegetation mapping was done by plant biologists in the field and spot-checked by senior biologists. The vegetation maps were digitized and converted to a GIS format. Acreage calculations by plant associations, dominant species and habitat type were done in GIS and GIS maps produced. Comparisons were made between the 1999 mapping and previous years' mapping from both the GIS data and the original data.

Comparison between years has shown that the surface area of total marsh and the distribution of marsh habitat types within the South Bay marshes is dynamic. New marsh formation (salt, brackish and freshwater marsh habitats) has occurred in both the Main Study Area (+202.02 acres) and the Reference Area (+52.78 acres). In the Main Study Area, new marsh formation occurred primarily in the Lower Reach (near the mouth of Coyote Creek) and Transition Reach. There has been little change in total marsh area in the Upper Reach during the past 10 years.

Overall, approximately 127 acres of salt marsh has converted to less saline habitat types within the Main Study Area between 1989 and 1999. The 127 acres of converted salt marsh represents 17% of the total salt marsh acreage present in the Main Study Area in 1989. Similarly, in the Reference Area approximately 16% (12 acres) of the total salt marsh area present in 1989 was converted to less saline habitat types during that same time period. The majority of this conversion has taken place since 1996. The majority of salt marsh habitat conversion during the past ten years is attributed to losses of pickleweed and cordgrass dominated associations and increases in alkali bulrush and peppergrass associations. In the Main Study Area, the majority of this conversion has occurred in the transition segments where nearly 100 acres of salt marsh habitat has become brackish marsh habitat during the past ten years.

There has been little net change in the total salt marsh area and in the proportion of salt marsh within the Main Study Area from 1989 to 1999. This apparent stability occurred because salt marsh habitat from conversion to other habitat types was balanced by increases in salt marsh habitat via new marsh formation. The entire study area has become less saline during the past ten years. Much of the change

has occurred since 1996; this was the first year that freshwater marsh habitats were mapped within the Main Study Area and the Reference Area.

Much of the conversion of salt marsh habitats within South San Francisco Bay are likely caused by large-scale influences that are affecting the entire system. This includes both anthropogenic and environmental factors. The impact from the WPCP can only be determined by a study that includes both physical and biological variables that could be influenced by freshwater flows. The ongoing collection of physical data (which began in August 1999) concurrent with this study will aid in determining the relative influences of environmental factors and anthropogenic factors to changes in marsh type.

INTRODUCTION

This report presents the results of vegetation mapping conducted in August and September 1999 in the South San Francisco Bay study area (Figure 1). The South Bay Marsh study area was subdivided into 24 mapping segments in the 1989 base study. Furthermore, the Alviso Slough Reference Reach was subdivided into 4 mapping segments. This was done to aid in determining specific locations of vegetation change.

For the purpose of data analysis and to better summarize vegetation change, the study area was organized into four reaches (Upper Reach segments, Transition Reach segments, Lower Reach segments and Alviso Slough Reference Reach) (Figure 2). The Upper, Transition and Lower Reach segments combined are referred to as the Main Study Area and are comprised of the original 24 mapping segments. The Alviso Slough Reference Reach is comprised of the 4 mapping segments established in the 1989 study.

The current study is a continuation of a long-term monitoring program designed to detect changes in the surface area and distribution of salt, brackish and freshwater marsh habitat types. An underlying objective of the long-term monitoring program is to evaluate the relative effect of freshwater discharge from the San Jose/Santa Clara Water Pollution Control Plant (WPCP) on the distribution and aerial extent of these habitat types. In an attempt to more accurately determine the relative contribution of the factors affecting marsh conversion, a study of edaphic characteristics (e.g. interstitial salinity, pH, and bulk density) and water levels throughout the study area was added to the monitoring program this year. Fieldwork commenced in August 1999. The results of soil characterization and water level monitoring as they relate to the 1999 distribution of marsh habitat types described herein will be presented in a separate report.

A baseline vegetation mapping study of the South San Francisco Bay study area was conducted in 1989 (H. T. Harvey & Associates 1990a). Subsequent mapping studies were conducted in 1991, 1994, 1995, 1996, 1997 and 1998 which documented changes in the distribution and aerial extent of salt, brackish and freshwater marsh (CH2MHill 1991; H.T. Harvey & Associates 1994, 1995, 1996, 1997, 1998). Comparison between years has shown that the distribution of plant associations within the South San Francisco Bay marshes is dynamic; the dominant plant species are adapted to respond rapidly to environmental changes. The 1998 study found that the overall marsh surface area (salt, brackish and freshwater marsh combined) increased by approximately 154 acres since 1989 in the Main Study Area. This increase was most likely due to sediment accretion and subsequent plant colonization by both brackish and salt marsh plant species. Approximately half of this increase (90 acres) occurred between 1997 and 1998 during the El Nino rainfall year.

Additionally, the 1998 study concluded that salt marsh area has decreased while brackish and freshwater marsh area has increased since 1989 within the Main Study Area (due to the conversion of both existing marsh and newly formed marsh). A nearly complete conversion of the Transition Reach segments from salt marsh to brackish marsh was observed. Between 1989 and 1998, a total of

approximately 126 acres of salt marsh (17% of the total) converted to primarily brackish marsh within the Main Study Area. Salt marsh conversion was primarily due to the successful establishment and vegetative reproduction of alkali bulrush (*Scirpus maritimus/robustus*) within mid-marsh areas that were formerly dominated by pickleweed (*Salicornia virginica*). A net decrease of 41 acres of salt marsh habitat (6% of the total) was also observed within the Main Study Area from 1989 to 1998. Due primarily to the formation of new cordgrass-dominated salt marsh, the net decrease in salt marsh area (41 acres) was considerably less than the surface area of salt marsh conversion (126 acres).

Pickleweed-dominated salt marsh is typically comprised of a diverse assemblage of mid-marsh salt tolerant species with pickleweed exhibiting the highest average percent cover. Pickleweed dominated salt marsh provides habitat for a unique assemblage of animal species including the federally and state-endangered salt marsh harvest mouse (*Reithrodontomys raviventris raviventris*). Therefore, it is important to determine the area of change annually as well as understand the factors responsible for the observed conversion of salt marsh to brackish marsh. Furthermore, it is important to understand to what extent this conversion is caused by natural, region-wide environmental change versus anthropogenic changes such as increases in freshwater discharge from the San Jose/Santa Clara WPCP and dry-weather releases from local reservoirs.

A number of variables have been shown to be important in controlling the distribution of plant species in coastal marshes. Interstitial soil salinity is one of the important variables correlated with vegetation change (Callaway and Sabraw 1994, Allison 1992, Callaway et al. 1989, Zedler 1983, 1986). For example, conversion of a pickleweed-dominated salt marsh to a cattail (*Typha dominguensis*)-dominated freshwater marsh was observed in the San Diego River marsh when reservoir discharges of freshwater were prolonged into summer; well beyond the normal rainy season (Zedler 1983). In this case study, prolonged freshwater discharge and increases in the depth and duration of inundation caused mortality of pickleweed and decreased soil salinities allowing cattail germination and growth (Zedler 1983).

However, numerous other factors have also been found to control marsh species composition including: depth and duration of flooding over the marsh surface (Webb and Mendelssohn 1996, Webb et al 1995, Pennings and Callaway 1992, Mendelssohn and McKee 1988), accumulation of phytotoxins such as hydrogen sulfide in marsh soils (Webb and Mendelssohn 1996, Webb et al. 1995, Koch and Mendelssohn 1989, DeLaune et al. 1983, King et al. 1982), interstitial nutrient concentrations (Koch et al. 1990, Bradley and Morris 1980, Koch and Mendelssohn 1989, Morris 1980) and soil mineral and organic matter content (Nyman et al. 1990, DeLaune et al. 1979). All of these variables can be affected by both natural and man-induced environment changes such as changes in precipitation, sea level and anthropogenic freshwater discharges. Increased flooding frequency due to sea level rise, for example, was implicated as the causal agent of plant association changes in a tidal marsh in the northeastern United States (Warren 1993).

Without empirical observations of the factors most likely to cause vegetation change within the study area (e.g. surface water and soil salinity, Coyote Creek flows) it has not been possible to definitively evaluate the relative effect of WPCP discharges on observed vegetation changes. However,

proportionate changes in overall marsh acreage and marsh types have occurred in both the Main Study Area and the Reference Reach (Alviso Slough). Assuming that the Reference Reach is not affected by WPCP discharges, this may indicate that large-scale, regional environmental changes are likely controlling the observed conversion of pickleweed-dominated salt marsh to brackish marsh in the Main Study Area. Specifically, annual precipitation exceeded the average from 1994 - 1998 (National Weather Service, Western Regional Climate Center) with the 1997 - 1998 El Nino rains (rainfall year 7/97 - 6/98) producing the third highest rainfall year on record since 1874 (28.89 inches, San Jose NWS substation at the civic center). Furthermore, increased freshwater flows from the Sacramento-San Joaquin Delta have been shown to significantly decrease salinities and increase the depth of flooding baywide (Dettinger et al. 1995, Peterson et al. 1995). Therefore, decreased salinity in the waters of the entire Bay and increased average tidal elevation associated with successive years of above average rainfall could be factors affecting changes in the South Bay marshes. Without concrete data regarding all of the factors above, the relative contribution of each is speculative.

The monitoring of surface water levels and salinity, soil bulk density, interstitial salinity and pH and flow rates from Coyote Creek and the Guadalupe River in combination with predicted tides for the South Bay and Delta Outflows will provide much of the information necessary to determine the relative effects of the discharge from the WPCP on observed vegetation changes. The addition of this data collection to the ongoing long-term vegetation monitoring will be presented in a future report.

Figure 1. Segment Locations

[figure 1.PDF](#)

Figure 2. Location of Four Reaches.

[figure 2.PDF](#)

SURVEY METHODS

STUDY AREA

For the purposes of data collection and analysis the study area was divided into 28 segments as defined in the 1989 study (H. T. Harvey & Associates 1990a; Figure 1). The study area was then divided into four reaches (Upper Reach segments, Transition Reach segments, Lower Reach segments, and Reference Reach) to provide a more easily comprehensible method of analyzing the data and presenting the results (Figure 2). The upper (approximately 440 acres), transition (approximately 391 acres), and Lower Reach (approximately 703 acres) segments, referred to as the Main Study Area are located within the Coyote Creek watershed and include Segments 1-5 and 8-26 (Figure 1). Segments 27-30 (Reference Area) are located along the Lower Reach segments of the Guadalupe River or Alviso Slough (Figure 1). These segments comprise a Reference Area (approximately 225 acres) for the purpose of documenting vegetation changes in a watershed not affected by the discharge of treated wastewater. The name and number of individual segments are listed in Table 1. These names were chosen for prominent geographic features displayed in Figure 1.

AERIAL PHOTOGRAPHY AND ORTHORECTIFICATION

The subconsultant responsible for aerial photography acquisition and digital imagery production, HJW, took color-infrared aerial (CIR) photographs of the entire study area. Color-infrared aerial photographs were taken on June 29, 1999. Photographs were taken from an altitude of 8500 feet using a 6-inch camera lens. The flight was scheduled during negative tidal elevation and 30 to 45 degree solar altitude.

This was the first year of this long-term study that the aerial photographs were orthorectified. The photographs were orthorectified to remove any distortion of the scale across the image caused by various factors including curvature of the earth's surface, topographic changes, and tilt of the camera lens. The use of orthorectified photographs adds greater accuracy to the estimation of polygon areas on the vegetation map. However, area calculations between previous year's data and the 1999 data will be influenced by this change in methodology.

The ortho processing procedure involved several consistent production steps, each including important inspections. First the film diapositive was scanned and thereby converted into a computer rasterized image. Scanning diapositives were made from the photography prior to any editing or other handling of the film. These diapositives were placed in individual sleeves to be kept free of dust, scratches, and any other blemishing agents. HJW maintains an environmentally controlled clean room for performing all photo scans to help eliminate airborne dust. The diapositives were scanned on a high precision Vexcel VS4000 scanner at the aperture of 25 microns. No pixels were resampled to convert to a finer resolution.

To correct an aerial photo for distortion caused by terrain; a digital terrain model (DTM) must be included in the ortho processing. HJW produced a DTM, not only capable of accurately generating the orthophotos, but sufficient for generating the digital elevation model (DEM) as well. Once scanned, HJW used OrthoView™ software to orthorectify the images and orient them into the California State Plane Coordinate System through the sensor orientation process. Control from the aerotriangulation and ground survey data from existing control points in HJW's database was used to tie the digital images to real world coordinates. The DTM collected from the stereo photography was used during the digital orthorectification process to adjust each image pixel into its correct position. HJW used a cubic convolution algorithm to perform the ortho processing. This technique provides a much more accurate solution than nearest neighbor methods.

Each image was visually checked and radiometrically enhanced if needed. Neighboring images were viewed and if problems were detected, they were feathered, or blended, along their edges to reduce radiometric differences. Where two adjoining images contain water, radiometric differences were not removed. Sun angles on water can result in severe tonal discontinuities that are quite labor intensive to repair. All digital orthophotographs were visually compared with the original unrectified image to verify radiometric accuracy.

VEGETATION ASSOCIATION MAPPING AND AREA CALCULATIONS

Field surveys and analysis of vegetation followed a protocol that began with mapping plant associations (comprised of either a single dominant individual plant or two dominant plants) onto clear acetate overlays that were placed directly over the digital images of the orthorectified CIR photos. These associations were subsequently assigned to one of three marsh types (i.e. salt marsh, brackish marsh or freshwater marsh) based upon the relative salinity tolerance of these species following the protocol established in the baseline study (H. T. Harvey & Associates 1990a). In order to facilitate comparison of results between monitoring years, vegetation associations were assigned to dominant species categories (as defined below). Dominant species categories, marsh types and vegetation associations for 1989, 1998 and 1999 are presented in Table 2.

Topographic features, marsh boundaries, and tentative vegetation associations (based on color signatures) were mapped in the office prior to field visits. Extensive ground-truthing of the preliminary mapping was then conducted during site visits to the entire Study Area conducted from 31 August to 22 September 1999. Marsh vegetation was observed primarily from areas directly adjacent to but not within the marshes in order to maintain consistency with the method employed in previous years. Marshes were, therefore, observed primarily from levee roadways, railroad beds, unimproved salt pond levees and Pacific Gas and Electric (PG&E) walkways.

When absolutely necessary, vegetation associations were verified by walking the marshes in areas that were not clearly visible from adjacent levees and upland areas

The field vegetation maps (acetate overlays) were then scanned and electronically digitized. The maps were then linked to the digital orthos images. Plant association acreages and color-coded figures for the entire Study Area were generated by GIS systems Microstation and ARCVIEW.

Access to the Study Area was obtained from the USFWS San Francisco Bay National Wildlife Refuge (Ms. Joy Albertson 510.792.0222; Special Use Permit Number 78109) and Cargill Salt Division, Newark, CA., (Mr. Chuck Taylor 510.797.1820; License Agreement 2001.009:98C).

Table 1. Segment Names and Numbers.

Segment Number	Segment Name
1	Calaveras Point, N
2	Calaveras Point, S
3	Calaveras Point, EI
4	Calaveras Point, EII
5	Albrae
8	Mowry Slough, Lower
9	Goose Point
10	Mud Slough, Lower W
11	Triangle
12	Coyote Creek, Mid S
13	Coyote Creek, Lower N
14	Coyote Creek, Lower N
15	Coyote Creek, Mid N
16	Coyote Creek, Upper
17	Warm Springs
18	Mud Slough, Upper
19	Mud Slough, Mid
20	Mud Slough, Lower E
21	Drawbridge
22	Strip
23	Knapp
24	Artesian Slough, Upper
25	Artesian Slough, Lower
26	Outfall
27	Alviso Slough, Gold St.
28	Alviso Slough, Upper
29	Alviso Slough, Middle
30	Alviso Slough, Lower

VEGETATION ASSOCIATION CATEGORIZATION METHODS

Any species that occurred as a dominant, co-dominant or sub-dominant in any portion of the study area was mapped. For the purposes of this study a dominant species had a percent cover of

Table 2. Dominant Species Categories, Marsh Type, and Vegetation Associations for 1989, 1998 and 1999.

DOMINANT SPECIES CATEGORY	HABITAT TYPE	VEGETATION ASSOCIATIONS		
		1989	1998	1999
Cordgrass	Salt	Cordgrass	Cordgrass Cordgrass/Sparscale Cordgrass/Alkali Bulrush Cordgrass/Pickleweed	Cordgrass Cordgrass/Alkali Bulrush Cordgrass/Pickleweed
Pickleweed	Salt	Pickleweed Pickleweed, Alkali Heath, Fat Hen	Pickleweed Pickleweed/Sparscale Pickleweed/Cordgrass Pickleweed/Peppergrass Pickleweed/Alkali Bulrush Pickleweed/Saltgrass	Pickleweed Pickleweed/Sparscale Pickleweed/Cordgrass Pickleweed/Peppergrass Pickleweed/Alkali Bulrush Pickleweed/Saltgrass Pickleweed/Gumplant Pickleweed/Alkali Heath
Pickleweed-Cordgrass Mix	Salt	Cordgrass, Pickleweed, with or without Alkali Heath	Pickleweed-Cordgrass Mix	Pickleweed-Cordgrass Mix
Pickleweed-Sparscale Mix	Salt	•	Pickleweed-Sparscale Mix	
Alkali Heath	Salt	•	Alkali Heath Alkali Heath/Alkali Bulrush	Alkali Heath
Gumplant	Salt	•	Gumplant Gumplant/Cordgrass Gumplant/Pickleweed Gumplant/Peppergrass Gumplant/Alkali Bulrush	Gumplant Gumplant/Cordgrass Gumplant/Pickleweed Gumplant/Peppergrass
Saltgrass	Salt	•	•	Saltgrass Saltgrass/Pickleweed
Jaumea	Salt	•	•	Jaumea Jaumea/Gumplant
Peripheral Halophytes	Salt	Fat Hen, Alkali Heath	Peripheral Halophytes Peripheral Halophytes/Peppergrass	Peripheral Halophytes
Alkali Bulrush	Brackish	Alkali Bulrush	Alkali Bulrush	Alkali Bulrush

DOMINANT SPECIES CATEGORY	HABITAT TYPE	VEGETATION ASSOCIATIONS		
		1989	1998	1999
			Alkali Bulrush/Pickleweed Alkali Bulrush/Peppergrass Alkali Bulrush/Sparscale Alkali Bulrush/Cordgrass Alkali Bulrush/California Bulrush Alkali Bulrush/Cattail Alkali Bulrush/Knotweed	Alkali Bulrush/Peppergrass Alkali Bulrush/Sparscale Alkali Bulrush/Cordgrass Alkali Bulrush/California Bulrush Alkali Bulrush/Cattail
Peppergrass	Brackish	Peppergrass	Peppergrass Peppergrass/Pickleweed Peppergrass/Alkali Bulrush Peppergrass/Sparscale Peppergrass/Peripheral Halophytes	Peppergrass Peppergrass/Pickleweed Peppergrass/Alkali Bulrush Peppergrass/Sparscale Peppergrass/Peripheral Halophytes Peppergrass/California Bulrush
Sparscale	Brackish	•	Sparscale Sparscale/Pickleweed Sparscale/Alkali Bulrush Sparscale/Peppergrass Sparscale/Gumplant Sparscale/Saltgrass	Sparscale Sparscale/Alkali Bulrush Sparscale/Peppergrass
California Bulrush	Fresh	•	California Bulrush California Bulrush/Knotweed California Bulrush/Cattail California Bulrush/Alkali Bulrush	California Bulrush California Bulrush/Knotweed California Bulrush/Cattail California Bulrush/Alkali Bulrush
Cattail	Fresh	•	Cattail Cattail/California Bulrush Cattail/Peppergrass Cattail/Alkali Bulrush	Cattail Cattail/California Bulrush Cattail/Alkali Bulrush
Knotweed	Fresh	•	•	Knotweed Knotweed/California Bulrush

- Not a Dominant Species Category in Analysis Year.

51-100%, co-dominant species have roughly equal percent coverage, and sub-dominant species have between 15 and 49 percent cover.

Each species was then assigned to a vegetation association comprised of one dominant, a dominant and subdominant, or two or more co-dominant species. The three types of vegetation associations are described below:

Dominant - An area that consists of one dominant species that comprises approximately 85-100% of the cover is named solely for that species alone, so that the vegetation association called Pickleweed consists of from 85-100% Pickleweed and of less than 15% of other unspecified species.

Dominant/sub-dominant - If one species comprises between approximately 51-85% of the cover in a particular area, and another species comprises 15-49% cover in that same area, then this is dominant/sub-dominant vegetation association. The association is named for both species, with the more abundant species listed first. The category called Pickleweed/Alkali bulrush could therefore consist of from 51-85% cover of Pickleweed and 15-49% cover of Alkali bulrush.

Co-dominant - Two co-dominant associations were identified: Pickleweed - Cordgrass (*Spartina foliosa*) Mix and Pickleweed - Spearscale (*Atriplex triangularis*) Mix. The species mixes represent an approximately equal amount of each species.

The upland species category consists of species not commonly found within salt marsh habitats. These include ruderal species such black mustard (*Brassica nigra*), ripgut grass (*Bromus diandrus*), bristly ox-tongue (*Picris echioides*), sweet fennel (*Foeniculum vulgare*), and coyote brush (*Baccharis pilularis*) as well as tree species such as California box elder (*Acer negundo ssp. californica*), California black walnut (*Juglans californica* var. *hindsii*) and Fremont cottonwood (*Populus fremontii*). The peripheral halophyte category consists of a patchwork of species that occur along salt marsh edges, adjacent to levees. This mixture includes pickleweed and various peripheral halophyte species such as alkali heath (*Frankenia salina*), Australian salt bush (*Atriplex semibaccata*) and slender-leaved iceplant (*Mesembryanthemum nodiflorum*).

Plant species associations were summarized into 15 dominant species categories (e.g. alkali bulrush/peppergrass association is an alkali bulrush dominant species category). These dominant species categories were then assigned to one of four habitat types: salt marsh, brackish marsh, freshwater marsh and upland. A number of assumptions about grouping dominant species into appropriate habitat types were made. These include:

- Relative salt tolerance of dominant plant species;
- Edaphic characteristics of the South Bay Marshes that may control plant species distribution;
- Historic relationships within this study, and;
- Relationships between dominant plant species and wildlife use.

Certain plant species for which salinity tolerance data is lacking (e.g. alkali bulrush, spearscale and peppergrass) were categorized into habitat types based on relative location in the marsh plain or known wildlife use. This assumption and the potential uncertainties related to assigning plant species to habitat type categories has been understood throughout the study period and was stated in the 1989 (baseline) study (H. T. Harvey & Associates 1990a). The habitat classification scheme first used in the baseline study is carried through to this study to maintain comparable data.

AREA COMPARISONS

Analysis of potential marsh conversion within the Main Study and Reference Areas involved a multi-step process that began at a total marsh area level and proceeded to a more specific, segment-level analysis. The first task involved comparing the relative acreage change in marsh type and dominant species categories between years. The current years results are compared to analysis year 1989. When a significant shift in marsh acreage occurred, the dominant species categories responsible for that shift were also identified.

In order to identify where significant acreage changes had occurred, the marsh was divided into four areas based upon segment location: Upper, Transition, Lower and Reference (Alviso Slough) (Figure 2). The Upper and Lower segments are upstream and downstream from the Transition segments, respectively. The Transition Segments include: 5, 9, 10, 11, 14 and 20. Upper Segments include: 12, 13, 15, 16, 17, 18, 19, 21, 24, 25 and 26. Lower Segments include: 1, 2, 3, 4, 8, 22 and 23. The Reference Area includes Segments 27, 28, 29 and 30.

A comparison of marsh habitat acreage data from all years (1989, 1991, 1994, 1996, 1997, 1998, and 1999) by location (reach) was also conducted to compare trends between reaches. The final step in the analysis overlaid the original data from previous years' mapping onto 1999 data to determine, with confidence, the location and size of change in marsh area and habitat type.

Dominant species and habitat maps were produced for each of the four segment locations. The maps were produced from an ArcView database and the full mapping for all segments by plant species association is available electronically.

RESULTS

GENERAL SPECIES DISTRIBUTION, DOMINANT SPECIES CATEGORY AND HABITAT ACREAGES FOR 1999

Main Study Area

Information is presented below by dominant species categories, which is a distillation of vegetation associations, and by habitat types. The spatial distribution of dominant plant species and habitat types are presented in Appendix A for each of the three segment locations within the Main Study Area (figure scales vary). Acreages of habitat types and associated dominant plant species for the Main Study Area are shown in Table 3 and Figure 3. The dominant plant species within the Main Study Area is alkali bulrush comprising just over 600 acres. The next prevalent species is pickleweed (Table 3). The total acreage of salt marsh habitat and brackish marsh habitat within the Main Study Area is nearly equal.

The Upper Reach segments consist primarily of brackish marsh associations dominated by either pure stands or mixtures of alkali bulrush and peppergrass (*Lepidium latifolium*). The Lower Reach segments (nearest San Francisco Bay) are comprised primarily of single-species stands or mixtures of the salt marsh plant species dominated by pickleweed and cordgrass. Although cordgrass and pickleweed are most abundant in the Lower Reach segments, both occur at low abundance even in the furthest upstream segments (although sometimes in patches too small to map). Conversely, peppergrass is most abundant in the Upper Reach segments, but is found throughout most of the Main Study Area (Appendix A).

Alkali bulrush occurs throughout the Main Study Area and is the dominant plant species of brackish marsh associations in South San Francisco Bay. Each year, alkali bulrush has been mapped further downstream (closer to San Francisco Bay). The furthest downstream patches of alkali bulrush were observed within Segments 3 and 22 of the Lower Reach segments area. Please note that in previous years the plant was identified as *Scirpus robustus* according to available conscription of the time (Munns and Keck 1968). The species, *S. maritimus*, was in fact not a recognized taxa in California's flora until 1993 after taxonomic changes were made as part of the development of the *Jepson Manual*; the plant was previously identified as *S. fluviatilis* with a described distribution restricted to lakes and rivers. In reality, both *S. robustus* and *S. maritimus* were observed and collected within the study area during the 1999 field surveys. The current and previous floras describe the plants as readily hybridizing with each other. In the current study we have chosen to describe the plant as *S. maritimus*. This species is very widespread and is known from the south and east coast regions of the United States.

The Transition Reach, intermediate to the furthest upstream and downstream reaches, supported significant amounts of both salt and brackish species, which sometimes occurred in mixed associations (both brackish and salt marsh plant species).

Table 3. Summary of Acreages of the Main Study Area by dominant species categories for each habitat type for 1999.

Dominant Species Category	1999
Salt Marsh Categories	
Cordgrass	102.05
Pickleweed	483.79
Pickleweed-Cordgrass Mix	87.67
Pickleweed-Spearscale Mix	0.00
Alkali Heath	1.40
Gumplant	37.12
Peripheral Halophytes	45.69
Misc Others	14.57
<i>Sub-Total</i>	772.29
Brackish Marsh Categories	
Alkali Bulrush	606.78
Peppergrass	162.49
Spearscale	8.11
Misc. Others	0.00
<i>Sub-Total</i>	777.38
Freshwater Marsh Categories	
California Bulrush	69.90
Cattail	4.24
Misc	0.04
<i>Sub-Total</i>	74.18
Upland	
<i>Sub-Total</i>	26.95
TOTAL	1650.80

Figure 3. Dominant Species Categories of the Main Study Area by Habitat Type, 1999.






Figure 4. Dominant Species Categories of the Reference Area by Habitat Type, 1999.



Reference Area (Alviso Slough)

The spatial distribution of dominant plant species and habitat types in the Reference Area are presented in Appendix A. The 1999 plant association acreages for Alviso Slough are presented in Table 4. Plant species within Alviso Slough have a general distribution similar to the Main Study Area in terms of a progression from freshwater to brackish and salt marsh species extending from upstream to the confluence with Coyote Creek. Acreage of habitat types and associated dominant plant species are shown in Figure 4. Alkali Bulrush is the dominant plant species within the Reference Area and brackish marsh habitat comprises almost twice the area of salt marsh habitat.

Brackish marsh associations occur throughout Alviso Slough. Patches of alkali bulrush were observed as far downstream as Segment 30. Aside from a trace amount, peppergrass was not observed further downstream than Segment 29. Freshwater marsh associations are concentrated in the upstream portions of the slough and salt marsh associations dominate the downstream areas.

Table 4. Summary of Acreages of the Reference Area (Alviso Slough) by dominant species categories for each habitat type for 1998 and 1999.

Dominant Species Category	1999
Salt Marsh Categories	
Cordgrass	15.33
Pickleweed	30.32
Peripheral Halophytes	12.67
Misc. Others	0.87
<i>Sub-Total</i>	59.19
Brackish Marsh Categories	
Alkali Bulrush	135.38
Peppergrass	22.51
Spearscale	0.06
Misc. Others	0.00
<i>Sub-Total</i>	157.95
Freshwater Marsh Categories	
California Bulrush	20.55
Cattail	11.38
Misc. Others	0.00
<i>Sub-Total</i>	31.93
Upland	
<i>Sub-Total</i>	18.16
TOTAL	267.23

TEMPORAL AND SPATIAL CHANGES IN MARSH HABITAT ACREAGES FROM 1989 THROUGH 1999

This comparison does not include data from segments 24, 25 and 26 of the Main Study Area and segment 27 of the Reference Area since they were not mapped in 1989. Additionally, the Reference Area was not mapped in 1994, therefore only data from the Main Study Area in 1994 is included in the evaluation.

New Marsh Formation (Salt, Brackish, and Freshwater Marsh Combined)

The surface area of marsh habitat has increased by 202.02 acres between 1989 and 1999 within the Main Study Area (Upper, Transition and Lower Reaches Combined) (Table 5). During the same period, 52.78 acres of new marsh has formed in the Reference Area (Table 6). This equates to a 15 % increase in marsh acreage in the Main Study Area and a 31 % increase in marsh acreage in the Reference Area between 1989 and 1999.

Marsh area remained relatively stable from 1989 to 1996 in the Main Study Area (Figure 5). The formation of new marsh habitat in the Main Study Area has occurred primarily between 1996 and 1999 in the Lower Reach and between 1996 and 1998 in the Transition Reach (Figure 5). Gains in marsh area were greatest in the Lower Reach (151.7 acres), while 49.5 acres of new marsh formation has occurred in the Transition Reach. The majority of new marsh formation has occurred in the Lower Reach along the north side of Coyote Creek, immediately upstream of Calaveras Point. Marsh area has increased steadily in the Lower Reach from 1996 through 1998 while the rate of increase appears to have slowed from 1998 to 1999 (Figure 5). In contrast, in the Transition Reach marsh area increased from 1996 through 1998 and remained comparable between 1998 and 1999 (Figure 5). Compared to the lower and Transition Reaches, the surface area of marsh in the Upper Reach has remained relatively stable throughout this 10 year study (Figure 5).

A trend of increasing marsh area is apparent from 1989 through 1999 in the Reference Area (Figure 5). The majority of this new marsh formation appears to have occurred between 1991 and 1996 and between 1997 and 1999.

Changes in Surface Area of Salt, Brackish, and Freshwater Marsh Habitats

Salt Marsh. Figures 6, 7 and 8 present the surface area of salt, brackish and freshwater marsh habitats by year and location (reach). Salt marsh area decreased in the Transition Reach from 1989 through 1999; the rate of decrease in salt marsh area was greatest between 1989 and 1994 (Figure 6). Conversely, salt marsh area increased in the Lower Reach from 1989 through 1999 with most of the increase occurring between 1996 and 1999. Much of this increase was likely due to new marsh formation along the north side of Coyote Creek downstream of Calaveras Point. Despite these changes, there has been little net change in salt marsh habitat area from 1989 to 1999 within the Main Study Area (Table 5). The net stability of salt marsh area within the Main Study Area was due to gains

from new marsh formation in the Lower Reach balancing losses in the Transition Reach (due to conversion) (Figure 6).

Table 5. Summary of Acreages of the Main Study Area by dominant species categories for each habitat type for 1989 and 1999*.

Dominant Species Category	1989	1999	Change
Salt Marsh Categories			
Cordgrass	70.70	102.05	31.35
Pickleweed	499.00	454.58	-44.42
Pickleweed-Cordgrass Mix	171.60	116.62	-54.98
Pickleweed-Spearscale Mix	0.00	0.00	0.00
Alkali Heath**	0.00	1.40	1.40
Gumplant**	0.00	37.12	37.12
Peripheral Halophytes	27.00	44.20	17.20
Misc Others	0.00	14.57	14.57
<i>Sub-Total</i>	768.30	770.54	2.24
Brackish Marsh Categories			
Alkali Bulrush	493.40	598.52	105.12
Peppergrass	70.90	144.82	73.92
Spearscale**	0.00	7.88	7.88
Misc. Others	0.00	0.00	0.00
<i>Sub-Total</i>	564.30	751.22	186.92
Freshwater Marsh Categories			
California Bulrush	0.00	10.69	10.69
Cattail	0.00	2.13	2.13
Misc	0.00	0.04	0.04
<i>Sub-Total</i>	0.00	12.86	12.86
TOTAL	1332.60	1534.62	202.02

* Comparison consists of segments 1-5, 8-23 only since segments 24-26 were not mapped in 1989.

** Not a dominant species category in 1989.


Table 6. Summary of Acreages of the Reference Area (Alviso Slough) by dominant species categories for each habitat type for 1989 and 1999.*

Dominant Species Category	1989	1999	Change
Salt Marsh Categories			
Cordgrass	27.90	15.33	-12.57
Pickleweed	46.80	30.32	-16.48
Peripheral Halophytes	2.60	12.66	10.06
Misc. Others	0.00	0.87	0.87
<i>Sub-Total</i>	<i>77.30</i>	<i>59.18</i>	<i>-18.12</i>
Brackish Marsh Categories			
Alkali Bulrush	77.10	127.99	50.89
Peppergrass	17.60	21.26	3.66
Spearscale**	0.00	0.06	0.06
Misc. Others	0.00	0.00	0.00
<i>Sub-Total</i>	<i>94.70</i>	<i>149.31</i>	<i>54.61</i>
Freshwater Marsh Categories			
California Bulrush	0.20	15.87	15.67
Cattail	0.00	0.62	0.62
Misc. Others	0.00	0.00	0.00
<i>Sub-Total</i>	<i>0.20</i>	<i>16.49</i>	<i>16.29</i>
TOTAL	172.20	224.98	52.78

* Comparison consists of segments 1-5, 8-23 only since segments 24-26 were not mapped in 1989.

** Not a dominant species category in 1989.

Figure 5. Total Marsh Acreage Comparison From 1989 to 1999 by Reach



*No data collected in 1994 within Reference Area.

A relatively large loss of salt marsh habitat has occurred in the Reference Area between 1989 and 1999 (Table 6). In contrast to the Main Study Area this loss was not compensated for by new salt marsh formation. Approximately 18.1 acres of salt marsh (24% of the total) has been lost during the study period and is comprised of losses in both pickleweed and cordgrass dominated categories. Similar to the pattern in the Transition Reach, the majority of salt marsh decline in the Reference Reach occurred early in the study period between 1991 and 1996 (Figure 6).

Brackish and Freshwater Marsh. Overall large gains in brackish marsh area have occurred in both the Main Study Area and in the Reference Area between 1989 and 1999 (Tables 5 and 6). During this period, brackish marsh increased by 186.92 acres (33% increase) and 54.61 acres (58% increase) in the Main Study and Reference Areas, respectively (Tables 5 and 6). This is due mostly to marsh conversion (from salt to brackish) but some new marsh has formed as brackish marsh. Furthermore, freshwater marsh has increased in the Main Study and Reference Area during the past 10 years (Tables 5 and 6).

In the Main Study Area, gains in brackish marsh have occurred in the lower and Transition Reaches while brackish marsh has decreased slightly in the Upper Reach (Figure 7). Expansion of brackish marsh area occurred primarily between 1997 and 1998 in the Upper Reach and from 1991 through 1998 in the Transition Reach (Figure 7). Similar to the Transition Reach, the Reference Area exhibited a trend of increasing brackish marsh area from 1991 through 1999 (Figure 7). Increases in freshwater marsh habitat have only occurred in the Upper Reach (Figure 8).

Figure 6. Salt Marsh Acreage Comparison From 1989 to 1999 by Reach.





Figure 7. Brackish Marsh Acreage Comparison From 1989 to 1999 by Reach



*No data collected in 1994 within Reference Area.




Figure 8. Freshwater Marsh Acreage Comparison From 1989 to 1999 by Reach

*No data collected in 1994 within Reference Area.

Temporal Changes in Proportional Area of Salt and Brackish Marsh Between the Main Study and Reference Areas

The proportion of salt marsh and brackish marsh area relative to total marsh area was compared between the Main Study and Reference Areas from 1989 through 1999 (Figures 9 and 10). This analysis was performed to control for the difference in size between the Main Study and Reference Areas. The percentage of salt marsh in the Main Study Area remained relatively stable from 1989 through 1997 with a slight decline between 1997 and 1998 (Figure 9). The decline in the percentage of salt marsh was greater in the Reference Area compared to the Main Study Area (Figure 9). The majority of this decline occurred between 1991 and 1996 in the Reference Area.

Similar to the pattern for salt marsh habitat, the percentage of brackish marsh has been relatively stable in the Main Study Area (Figure 10). Within the Main Study Area, slight increases in the proportion of brackish marsh were observed between 1989 and 1991 and between 1997 and 1998 (Figure 10). A larger increase in the percentage of brackish marsh was observed in the Reference Area compared to the Main Study Area (Figure 10). This increase in the proportion of brackish marsh area to total marsh area in the Reference Area occurred primarily between 1991 and 1996 (Figure 10) during the same time that the percentage of salt marsh declined (Figure 9).

Figure 9. Temporal Comparison of the Proportion of Salt Marsh Area Between the Main Study and Reference Areas





Figure 10. Temporal Comparison of the Proportion of Brackish Marsh Area Between the Main Study and Reference Areas



*No data collected in 1994 within Reference Area.

Table 7. Detailed Evaluation of Changes in Acreage for Segment Locations by Habitat Type and for Total Marsh, 1989 to 1999.

Segment Location	Change in SM acreage	Change in BM acreage	Change in FM acreage	Change in TM acreage	Evaluation Results
Lower	93.91	62.27	0	156.61	Increase in TM acreage is due to approximately 107 acres of new marsh formation primarily in segments 3 and 4. This increased marsh area has formed as both SM and BM habitat which, in part, accounts for the increase in these habitat types. Approximately 46 acres of SM has converted to BM during the past ten years. The remaining change is due to mapping error. Net SM Conversion: 46 acres.
Transition	-94.5	133.4	.02	39.25	Approximately 8 acres of the TM acreage increase is due to new marsh formation. Approximately 76 acres of salt marsh has converted to BM, primarily in segments 5, 9, 10, and 11. The remaining change is due to mapping error. Net SM Conversion: 76 acres.
Upper	2.97	-8.7	12.85	30.42	The increase in TM acreage is due to approximately 9 acres of new marsh formation primarily in FM habitat. Approximately 5 acres of SM has converted to BM while approximately 14 acres of BM has converted to FM. The remaining change is due to mapping error. Net SM Conversion: 5 acres.
Reference	-18.12	54.65	16.23	58.50	Approximately 13 acres of new marsh formation has occurred during the past ten years. This marsh has formed primarily as BM and FM habitats. Approximately 12 acres of SM has converted to BM. The remaining change is due to mapping error. Net SM Conversion: 12 acres.

Total SM Conversion Within Main Study Area= 127 acres (17% of total SM acreage in 1989).

Total SM Conversion Within Reference Area = 12 acres (16% of total SM acreage in 1989).

SM = Salt Marsh Habitat

BM = Brackish Marsh Habitat

FM = Freshwater Marsh Habitat

TM = Total Marsh Area

Habitat Type Conversion

Detailed comparisons of the original overlays were conducted to isolate specific areas of major habitat change. Table 7 provides a summary of the segment locations and detailed explanation of significant shifts in acreage by marsh type and/or total marsh area from 1989 to 1999. A total of 127 acres of salt marsh habitat (17% of the total) has converted to brackish marsh habitat from 1989 to 1999 in the Main Study Area. During the same period, 12 acres (16% of the total) of salt marsh habitat has converted to brackish marsh in the Reference Area. The remaining change indicated by the GIS data could not be accounted for through detailed analysis of the original data. That difference can be accounted for in minor changes within segments and mapping associated errors.

DISCUSSION

There has been an apparent increase of approximately 226.3 acres of overall mapped area (202 acres of new marsh formation) since 1989 in the Main Study Area. The majority of this increase is apparently due to sediment accretion along slough and river channels and subsequent vegetation colonization to form new marsh area. Some portion of the overall increase in marsh area can be attributed to differences in mapping methodology between years and errors associated with previous years' photography that is not orthorectified.

The majority of new marsh formation continues to occur at Calaveras Point located at the mouth of Coyote Creek, which has increased 16 acres during the past year and has grown in size from 173 acres in 1989 to 279 acres in 1999 (a 62% increase). It appears that substantial sedimentation along Coyote Creek has raised the elevations to a level that will support the growth of emergent plant species. This newly formed mud flat continues to be colonized by alkali bulrush and a mixture of cordgrass and annual pickleweed (*Salicornia europaea*).

Within the Main Study Area from 1989 to 1999, losses in salt marsh habitat from conversion to other habitat types were balanced by increases in salt marsh habitat via new marsh formation. The majority of salt marsh habitat conversion during the past ten years is attributed to losses of pickleweed and cordgrass dominated associations and increases in alkali bulrush and peppergrass associations. Furthermore, the majority of this conversion has occurred in the transition segments where nearly 100 acres of salt marsh habitat has become brackish marsh habitat during the past ten years. Conversely during the past year approximately 2 acres of salt marsh conversion has occurred within the lower reach segments. Much of the conversion that has occurred during the past two years has been in an area disjoint from brackish marsh habitat within the transition segments; prior to 1998 most of the conversion was occurring farther upstream.

Historically, the channel-side vegetation in the transition segments may have been dominated by brackish (alkali bulrush) and freshwater species (tules) (SFEI 1999). Salt marsh habitat dominated by pickleweed and salt grass occurred inland of the channel-side vegetation (SFEI 1999). Those areas that were historically salt marsh have likely been converted to salt ponds. Many of the existing marshes, located between the levees of the salt ponds and the channels, have formed more recently. The remnant historical marshes are likely the present day channel-side brackish marshes that were at one time small patches of low salinity marshes within a larger matrix of salt marsh habitat (SFEI 1999).

The entire study area is becoming less saline. No freshwater marsh habitat was mapped prior to 1996 in the Main Study Area or Alviso Slough (except in Segments 26 and 27 which are not part of the 10-year analysis) but now accounts for 29 acres within the study area. Newly forming freshwater marsh habitat in both the Reference Area and the Main Study Area indicates that freshwater influences are affecting all marshes of South San Francisco Bay. Additionally, the net salt marsh acreage within the Main Study Area has been relatively stable during this period of increased freshwater impacts. The stability in salt marsh acreage during a period when salt marsh conversion is predominate is due to a

simultaneous increase in new salt marsh via marsh formation and a loss of salt marsh habitat because of conversion to brackish marsh.

The relative change in habitat types through time is similar between the Main Study Area and Reference Area although the rate of new marsh formation in the Main Study Area has exceeded that of the Reference Area. This indicates that much of the conversion of salt marsh habitats within the South San Francisco Bay area is likely driven by large-scale influences that are affecting the entire system. This includes both environmental (e.g. precipitation and delta outflows) and anthropogenic factors (e.g. WPCP discharge and dry-weather releases from local reservoirs).

The impact from the WPCP plant can only be determined from a study that includes both physical and biological variables that could be influenced by the freshwater flows. This study has been expanded to include the collection of surface water and porewater data and will be presented in a report at a later date. Furthermore, improvements in the base materials used to conduct the vegetation mapping will allow future datasets to be more accurate and better reflect real changes in the South Bay Marshes. As the physical data currently being collected is combined with the information presented in this report, the relative influences of environmental factors and anthropogenic factors can be better understood.

REFERENCES

- Allison, S. K. 1992. The influence of rainfall variability on the species composition of a northern California salt marsh plant assemblage. *Vegetation* 101: 145-160.
- Bradley, P.M. and J.T. Morris. 1980. Influence of oxygen and sulfide concentration on nitrogen uptake kinetics in *Spartina alterniflora*. *Ecology*. 71:282-288.
- Callaway, R. M., Jones, S., Ferren, W. R., Jr., and A. Parikh. 1989. Ecology of a mediterranean-climate estuarine wetland at Carpinteria, California: plant distributions and soil salinity in the upper marsh.
- Callaway, R. M. and C. S. Sabraw. 1994. Effects of variable precipitation on the structure and diversity of a California salt marsh community. *Journal of Vegetation Science* 5: 433-438.
- CH2M Hill. 1991. Marsh Plant Associations of South San Francisco Bay: 1991 Comparative Study. 16 pp.
- DeLaune, R. D., Buresh, R. J., and W. H. Patrick, Jr. 1979. Relationship of soil properties to standing crop biomass of *Spartina alterniflora* in a Louisiana marsh. *Estuarine and Coastal Marine Science* 8:477-487.
- DeLaune, R. D., Smith, C. J. And W. H. Patrick, Jr., 1983. Relation of marsh elevation, redox potential and sulfide to *Spartina alterniflora* productivity. *Soil Science Society of America Journal* 47:930-935.
- Dettinger, Michael, Richard Smith, Daniel Cayan, David Peterson and Reginal Uncles. 1995. Hydroclimatology of San Francisco Bay Freshwater Inflows and Salinity, U. S. Geological Survey Fact Sheet, 5 p.
- Harvey, H. T. & Associates. 1984. South Bay Dischargers Authority Comparative Study. No. 156-02.
- Harvey, H. T. & Associates. 1990a. Marsh Plant Associations of the South Bay: 1989 Baseline Study. No. 477-05.
- Harvey, H. T. & Associates. 1990b. Plant Association Acreage Changes in South Bay: 1985-1989. No. 477-10.
- Harvey, H. T. & Associates. 1994. Marsh Plant Associations of South San Francisco Bay: 1994 Comparative Study. No. 477-14.

- Harvey, H. T. & Associates. 1995. Marsh Plant Associations of Artesian Slough and Transition Zone, South San Francisco Bay: Appendix to the 1994 Comparative Study. No. 477-14.
- Harvey, H. T. & Associates. 1997. Marsh Plant Associations of South San Francisco Bay: 1996 Comparative Study Including Alviso Slough. No. 477-18.
- Harvey, H. T. & Associates. 1998. Marsh Plant Associations of South San Francisco Bay: 1997 Comparative Study. No. 447-19
- King, G., Klug, M. J., Wiegert, R. G., and A. G. Chalmers. 1982. Relation of soil water movement and sulfide concentration to *Spartina alterniflora* production in a Georgia salt marsh. *Science* 218:61-63.
- Koch, M. S. and I. A. Mendelsohn. 1989. Sulfide as a soil phytotoxin: Differential responses in two salt marsh species. *Journal of Ecology* 77:565-578.
- Koch, M. S., Mendelsohn, I. A., and K. L. McKee. 1990. Mechanisms for the hydrogen sulfide-induced growth limitation in wetland macrophytes. *Limnology and Oceanography* 35:395-408.
- Mendelsohn, I. A., and K. L. McKee. 1988. *Spartina alterniflora* die-back in Louisiana: Time-course investigation of soil waterlogging effects. *Journal of Ecology* 76:509-521.
- Morris, J. T. 1980. The nitrogen uptake kinetics of *Spartina alterniflora* in culture. *Ecology* 61:1114-1121.
- Nyman, J. A., DeLaune, R. D. and W. H. Patrick, Jr. 1990. Wetland soil formation in the rapidly subsiding Mississippi River deltaic plain: Mineral and organic matter relationships. *Estuarine, Coastal and Shelf Science* 31:57-69.
- Pennings, S. C. And R. M. Callaway. 1992. Salt marsh plant zonation: The relative importance of competition and physical factors. *Ecology* 72:681-690.
- Peterson, D.H., Cayan, D.R., Dettinger, M.D., Hager, S.E., Schemel, L.E., Smith, R.E., DiLeo, J., and Uncles, R.E., 1995, Seasonal/yearly variation in San Francisco Bay: U.S. Geological Survey Fact Sheet, 8 p.
- San Francisco Estuary Institute. 1999. Conceptual Models of Freshwater Influences on Tidal Marsh Form and Function, with an Historical Perspective. 61p.
- Warren, R. S. and W. A. Niering. 1993. Vegetation changes on a northeast tidal marsh: interaction of sea-level rise and marsh accretion. *Ecology* 74:96-103.
- Webb, E. C., Mendelsohn, I.A., and B. J. Wilsey. 1995. Causes for vegetation dieback in a Louisiana salt marsh: A bioassay approach. *Aquatic Botany* 51:281-289.

Webb, E. C. and I. A. Mendelssohn. 1996. Factors affecting vegetation dieback of an oligohaline marsh in coastal Louisiana: Field manipulation of salinity and submergence. *American Journal of Botany* 83: 1429-1434.

Zedler, J. B. 1983. Freshwater impacts in normally hypersaline marshes. *Estuaries* 6:346-355.

Zedler, J.B. and P.A. Beare. 1986. Temporal variability of salt marsh vegetation: the role of low-salinity gaps and environmental stress. *In* (eds.) *Estuarine Variability*. Academic Press, Inc.

APPENDIX A
SOUTH BAY MARSHES:
1999 VEGETATION MAPS

[1999maps.pdf](#)

APPENDIX B
VEGETATION MATRICES

**Table B1. Acreage Summary of Segment 1 for 1989, 1994/1995, 1996, 1997, 1998 and 1999.
DOMINANT SPECIES CATEGORY**

<u>Salt marsh Vegetation</u>	<u>Year</u>					
	1989	1994/1995	1996	1997	1998	1999
Pickleweed	0.0	19.2	27.2	18.6	12.2	12.6
Cordgrass	8.2	1.4	3.4	2.8	9.7	1.94
Pickleweed-Cordgrass Mix	14.1	0.0	0.0	1.3	0.8	0.7
Pickleweed-Spearscale Mix	0.0	0.0	0.0	0.0	0.0	0.0
Alkali Heath	0.0	0.0	0.0	0.0	0.0	0.0
Gumplant	0.0	0.0	0.0	0.0	0.2	0.1
Peripheral Halophytes	2.6	1.5	1.7	0.0	1.4	1.43
Total Saline Dominant Species:	24.9	22.1	32.3	22.7	24.3	16.8
<u>Brackish Marsh Vegetation</u>						
Alkali Bulrush	0.0	0.0	0.0	0.0	0.0	0.0
Peppergrass	0.0	0.0	0.0	0.0	0.0	0.0
Spearscale	0.0	0.0	0.0	0.0	0.0	0.0
Total Brackish Dominant Species:	0.0	0.0	0.0	0.0	0.0	0.0
<u>Freshwater Marsh Vegetation</u>						
California Bulrush	0.0	0.0	0.0	0.0	0.0	0.0
Cattail	0.0	0.0	0.0	0.0	0.0	0.0
Total Freshwater Dominant Species:	0.0	0.0	0.0	0.0	0.0	0.0
Total Segment Acreage	24.9	22.1	32.3	23.3	26.5	27.1

**Table B2. Acreage Summary of Segment 2 for 1989, 1994/1995, 1996, 1997, 1998 and 1999.
DOMINANT SPECIES CATEGORY**

<u>Salt Marsh Vegetation</u>	<u>Year</u>					
	1989	1994/1995	1996	1997	1998	1999
Pickleweed	26.0	35.5	32.9	32.4	19.0	36.2
Cordgrass	13.3	2.3	2.6	3.8	10.5	3.1
Pickleweed-Cordgrass Mix	0.0	0.0	0.0	1.8	0.0	0.0
Pickleweed-Spearscale Mix	0.0	0.0	0.0	0.0	0.0	0.0
Alkali Heath	0.0	0.0	0.0	0.0	0.0	0.0
Gumplant	0.0	0.0	0.0	0.2	0.2	1.4
Peripheral Halophytes	3.6	2.3	1.6	0.7	3.0	2.2
Total Saline Dominant Species:	42.9	40.1	37.1	38.9	32.7	42.9
<u>Brackish Marsh Vegetation</u>						
Alkali Bulrush	0.0	0.0	0.0	0.0	0.0	0.0
Peppergrass	0.0	0.0	0.0	0.0	0.0	0.0
Spearscale	0.0	0.0	0.0	0.4	7.5	0.0
Total Brackish Dominant Species:	0.0	0.0	0.0	0.4	7.5	0.0
<u>Freshwater Marsh Vegetation</u>						
California Bulrush	0.0	0.0	0.0	0.0	0.0	0.0
Cattail	0.0	0.0	0.0	0.0	0.0	0.0
Total Freshwater Dominant Species:	0.0	0.0	0.0	0.0	0.0	0.0

Total Segment Acreage	42.9	40.1	37.1	39.8	41.2	42.9
------------------------------	------	------	------	------	------	------

Table B3. Acreage Summary of Segment 3 for 1989, 1994/1995, 1996, 1997, 1998 and 1999.

DOMINANT SPECIES CATEGORY

	<u>Year</u>					
	1989	1994/1995	1996	1997	1998	1999
<u>Salt Marsh Vegetation</u>						
Pickleweed	97.8	114.7	79.3	95.1	98.7	118.3
Cordgrass	4.1	3.4	2.9	86.6	104.6	15.9
Pickleweed-Cordgrass Mix	68.6	69.9	98.8	36.0	0.0	83.3
Pickleweed-Spearscale Mix	0.0	0.0	0.0	0.0	0.1	0.0
Alkali Heath	0.0	0.0	0.0	0.0	0.0	0.0
Gumplant	0.0	0.0	2.7	6.9	2.2	7.4
Peripheral Halophytes	2.4	2.6	1.1	1.0	2.2	1.0
Total Saline Dominant Species:	172.9	190.6	184.8	225.6	207.9	225.9
<u>Brackish Marsh Vegetation</u>						
Alkali Bulrush	0.0	0.0	0.1	0.0	49.2	50.8
Peppergrass	0.0	1.1	1.2	1.6	1.8	1.8
Spearscale	0.0	0.0	0.0	0.2	2.4	0.0
Total Brackish Dominant Species:	0.0	1.1	1.3	1.8	53.4	52.6
<u>Freshwater Marsh Vegetation</u>						
California Bulrush	0.0	0.0	0.0	0.0	0.0	0.0
Cattail	0.0	0.0	0.0	0.0	0.0	0.0
Total Freshwater Dominant Species:	0.0	0.0	0.0	0.0	0.0	0.0
Total Segment Acreage	172.9	191.7	212.3	227.6	262.1	278.5

Table B4. Acreage Summary of Segment 4 for 1989, 1994/1995, 1996, 1997, 1998 and 1999.

DOMINANT SPECIES CATEGORY

	<u>Year</u>					
	1989	1994/1995	1996	1997	1998	1998
<u>Salt Marsh Vegetation</u>						
Pickleweed	36.4	43.9	46.9	50.1	49.8	47.6
Cordgrass	6.1	6.2	4.1	5.6	12.9	17.1
Pickleweed-Cordgrass Mix	7.0	3.4	6.2	7.2	0.1	0.0
Pickleweed-Spearscale Mix	0.0	0.0	0.0	0.0	0.0	0.0
Alkali Heath	0.0	0.0	0.0	0.2	0.0	0.0
Gumplant	0.0	0.0	0.0	0.1	0.1	0.1
Peripheral Halophytes	0.4	2.4	1.5	0.9	1.7	1.7
Total Saline Dominant Species:	49.9	55.9	58.7	64.0	64.6	66.5
<u>Brackish Marsh Vegetation</u>						
Alkali Bulrush	0.0	0.0	0.0	0.0	4.8	6.2
Peppergrass	0.0	0.1	0.1	0.1	0.1	0.2
Spearscale	0.0	0.0	0.0	0.0	0.1	0.0
Total Brackish Dominant Species:	0.0	0.1	0.1	0.1	5.0	6.4
<u>Freshwater Marsh Vegetation</u>						
California Bulrush	0.0	0.0	0.0	0.0	0.0	0.0
Cattail	0.0	0.0	0.0	0.0	0.0	0.0

Total Freshwater Dominant Species:	0.0	0.0	0.0	0.0	0.0	0.0
Total Segment Acreage	49.9	56.0	58.8	64.0	70.0	72.9

Table B5. Acreage Summary of Segment 5 for 1989, 1994/1995, 1996, 1997, 1998 and 1999.
DOMINANT SPECIES CATEGORY

	<u>Year</u>					
	1989	1994/1995	1996	1997	1998	1999
<u>Salt Marsh Vegetation</u>						
Pickleweed	60.5	62.3	30.5	36.6	34.4	41.6
Cordgrass	0.5	2.1	2.7	2.6	3.6	2.3
Pickleweed-Cordgrass Mix	0.0	0.0	0.0	0.0	0.0	1.1
Pickleweed-Spearscale Mix	0.0	0.0	18.9	7.9	2.2	0.0
Alkali Heath	0.0	0.0	0.0	0.2	0.1	0.4
Gumplant	0.0	0.0	0.0	0.1	0.0	0.3
Peripheral Halophytes	1.2	0.5	1.0	2.8	3.2	6.6
Total Saline Dominant Species:	62.2	64.9	53.1	50.2	43.5	52.3
<u>Brackish Marsh Vegetation</u>						
Alkali Bulrush	25.1	19.2	27.3	32.1	34.7	32.0
Peppergrass	0.0	1.4	2.4	4.0	3.4	7.5
Spearscale	0.0	0.0	0.0	3.7	13.6	0.1
Total Brackish Dominant Species:	25.1	20.6	29.7	39.8	51.7	39.6
<u>Freshwater Marsh Vegetation</u>						
California Bulrush	0.0	0.0	0.0	0.0	0.0	0.0
Cattail	0.0	0.0	0.0	0.0	0.0	0.0
Total Freshwater Dominant Species:	0.0	0.0	0.0	0.0	0.0	0.0
Total Segment Acreage	87.3	85.5	82.8	90.0	95.5	92.4

Table B6. Acreage Summary of Segment 8 for 1989, 1994/1995, 1996, 1997, 1998 and 1999.
DOMINANT SPECIES CATEGORY

	<u>Year</u>					
	1989	1994/1995	1996	1997	1998	1999
<u>Salt Marsh Vegetation</u>						
Pickleweed	188.0	204.9	151.8	149.4	101.0	171.1
Cordgrass	13.1	11.7	10.2	22.5	98.0	32.5
Pickleweed-Cordgrass Mix	23.0	0.0	49.0	25.7	0.0	0.0
Pickleweed-Spearscale Mix	0.0	0.0	0.0	0.0	0.0	6.6
Alkali Heath	0.0	0.0	0.0	0.1	0.0	0.3
Gumplant	0.0	0.0	0.0	23.8	25.7	27.5
Peripheral Halophytes	7.1	10.0	7.8	6.0	10.1	7.7
Total Saline Dominant Species:	231.2	226.6	218.8	227.5	234.8	245.7
<u>Brackish Marsh Vegetation</u>						
Alkali Bulrush	0.0	0.0	0.0	0.0	0.0	0.0
Peppergrass	0.0	0.0	0.0	0.0	0.0	0.0
Spearscale	0.0	0.0	0.0	0.1	0.0	0.0
Total Brackish Dominant Species:	0.0	0.0	0.0	0.1	0.0	0.0
<u>Freshwater Marsh Vegetation</u>						
California Bulrush	0.0	0.0	0.0	0.0	0.0	0.0

Cattail	0.0	0.0	0.0	0.0	0.0	0.0
Total Freshwater Dominant Species:	0.0	0.0	0.0	0.0	0.0	0.0
Total Segment Acreage	231.2	226.6	215.3	228.5	239.1	248.7

Table B7. Acreage Summary of Segment 9 for 1989, 1994/1995, 1996, 1997, 1998 and 1999.

DOMINANT SPECIES CATEGORY

	Year					
	1989	1994/1995	1996	1997	1998	1999
<u>Salt Marsh Vegetation</u>						
Pickleweed	3.6	32.4	15.4	10.0	3.5	6.0
Cordgrass	3.7	8.9	3.9	6.6	7.3	4.7
Pickleweed-Cordgrass Mix	42.6	0.0	0.0	0.1	0.0	0.2
Pickleweed-Spearscale Mix	0.0	0.0	0.3	0.5	0.0	0.0
Alkali Heath	0.0	0.0	0.0	0.0	0.0	0.2
Gumplant	0.0	0.0	0.0	0.0	0.0	0.3
Peripheral Halophytes	1.2	0.0	1.3	2.0	3.3	1.2
Total Saline Dominant Species:	51.1	41.3	20.9	19.2	14.1	12.6
<u>Brackish Marsh Vegetation</u>						
Alkali Bulrush	16.3	22.2	44.1	50.4	67.0	60.2
Peppergrass	0.6	1.3	1.2	1.7	1.4	4.3
Spearscale	0.0	0.0	0.0	1.5	1.9	3.0
Total Brackish Dominant Species:	16.9	23.5	45.3	53.6	70.2	67.5
<u>Freshwater Marsh Vegetation</u>						
California Bulrush	0.0	0.0	0.0	0.0	0.0	0.0
Cattail	0.0	0.0	0.0	0.0	0.0	0.0
Total Freshwater Dominant Species:	0.0	0.0	0.0	0.0	0.0	0.0
Total Segment Acreage	68.0	64.8	66.2	72.8	84.8	80.2

Table B8. Acreage Summary of Segment 10 for 1989, 1994/1995, 1996, 1997, 1998 and 1999.

DOMINANT SPECIES CATEGORY

	Year					
	1989	1994/1995	1996	1997	1998	1999
<u>Salt Marsh Vegetation</u>						
Pickleweed	21.7	21.2	10.7	10.4	8.3	8.0
Cordgrass	6.8	11.0	8.4	8.3	5.0	3.6
Pickleweed-Cordgrass Mix	0.0	0.0	0.0	0.0	0.0	0.1
Pickleweed-Spearscale Mix	0.0	0.0	0.0	0.0	0.0	0.0
Alkali Heath	0.0	0.0	0.0	0.0	0.0	0.1
Gumplant	0.0	0.0	0.0	0.0	0.0	0.0
Peripheral Halophytes	0.7	0.1	0.6	0.6	1.6	0.2
Total Saline Dominant Species:	29.2	32.3	19.7	19.3	14.9	12.0
<u>Brackish Marsh Vegetation</u>						
Alkali Bulrush	9.7	5.8	19.7	24.3	37.1	30.7
Peppergrass	2.6	1.7	1.6	2.7	1.7	6.3
Spearscale	0.0	0.0	0.0	0.0	0.1	0.0
Total Brackish Dominant Species:	12.3	7.5	21.3	27.0	38.9	37.0
<u>Freshwater Marsh Vegetation</u>						

California Bulrush	0.0	0.0	0.0	0.0	0.0	0.0
Cattail	0.0	0.0	0.0	0.0	0.0	0.0
Total Freshwater Dominant Species:	0.0	0.0	0.0	0.0	0.0	0.0
Total Segment Acreage	41.5	39.8	41.0	46.3	53.8	49.0

Table B9. Acreage Summary of Segment 11 for 1989, 1994/1995, 1996, 1997, 1998 and 1999.
DOMINANT SPECIES CATEGORY

	<u>Year</u>					
	1989	1994/1995	1996	1997	1998	1999
<u>Salt Marsh Vegetation</u>						
Pickleweed	2.0	22.4	3.8	3.9	1.7	1.8
Cordgrass	0.0	1.6	1.1	1.1	1.6	2.3
Pickleweed-Cordgrass Mix	16.3	0.0	0.0	0.0	0.0	0.0
Pickleweed-Spearscale Mix	0.0	0.0	0.1	0.1	0.1	0.0
Alkali Heath	0.0	0.0	0.0	0.0	0.0	0.0
Gumplant	0.0	0.0	0.0	0.0	0.0	0.0
Peripheral Halophytes	1.0	0.0	0.4	1.1	1.5	1.2
Total Saline Dominant Species:	19.3	24.0	5.4	6.4	5.0	5.3
<u>Brackish Marsh Vegetation</u>						
Alkali Bulrush	51.2	48.8	63.4	64.4	68.5	68.6
Peppergrass	7.7	5.6	6.2	6.4	5.5	8.2
Spearscale	0.0	0.0	0.0	1.2	1.1	0.4
Total Brackish Dominant Species:	58.9	54.4	69.6	72.0	75.1	77.2
<u>Freshwater Marsh Vegetation</u>						
California Bulrush	0.0	0.0	0.0	0.0	0.0	0.0
Cattail	0.0	0.0	0.0	0.0	0.0	0.0
Total Freshwater Dominant Species:	0.0	0.0	0.0	0.0	0.0	0.0
Total Segment Acreage	78.2	78.4	75.1	78.3	80.7	82.6

Table B10. Acreage Summary of Segment 12 for 1989, 1994/1995, 1996, 1997, 1998 and 1999.
DOMINANT SPECIES CATEGORY

	<u>Year</u>					
	1989	1994/1995	1996	1997	1998	1999
<u>Salt Marsh Vegetation</u>						
Pickleweed	0.1	2.8	0.6	2.0	0.7	0.5
Cordgrass	0.0	2.2	1.1	1.1	0.7	1.4
Pickleweed-Cordgrass Mix	0.0	0.0	0.0	0.0	0.0	0.0
Pickleweed-Spearscale Mix	0.0	0.0	0.4	0.0	0.0	0.0
Alkali Heath	0.0	0.0	0.0	0.1	0.1	0.0
Gumplant	0.0	0.0	0.0	0.0	0.0	0.0
Peripheral Halophytes	0.6	0.0	1.7	1.1	10.2	2.2
Total Saline Dominant Species:	0.7	5.0	3.8	4.3	11.7	4.1
<u>Brackish Marsh Vegetation</u>						
Alkali Bulrush	28.2	21.2	25.4	24.1	19.0	24.2
Peppergrass	13.5	17.5	13.4	14.5	9.9	18.4
Spearscale	0.0	0.0	0.0	0.5	1.7	0.0
Total Brackish Dominant Species:	41.7	38.7	38.8	39.0	30.6	42.6

<u>Freshwater Marsh Vegetation</u>						
California Bulrush	0.0	0.0	0.1	0.2	0.2	0.2
Cattail	0.0	0.0	0.0	0.0	0.0	0.0
Total Freshwater Dominant Species:	0.0	0.0	0.1	0.2	0.2	0.2
Total Segment Acreage	42.4	43.7	43.1	43.5	44.5	47.4

Table B11. Acreage Summary of Segment 13 for 1989, 1994/1995, 1996, 1997, 1998 and 1999.
DOMINANT SPECIES CATEGORY

<u>Salt Marsh Vegetation</u>	<u>Year</u>					
	1989	1994/1995	1996	1997	1998	1999
Pickleweed	0.0	0.4	0.8	1.5	0.5	0.4
Cordgrass	0.0	0.4	0.0	0.2	0.0	0.0
Pickleweed-Cordgrass Mix	0.0	0.0	0.0	0.0	0.0	0.0
Pickleweed-Spearscale Mix	0.0	0.0	0.0	0.0	0.0	0.0
Alkali Heath	0.0	0.0	0.0	0.0	0.0	0.0
Gumplant	0.0	0.0	0.0	0.0	0.0	0.0
Peripheral Halophytes	0.7	0.0	11.9	7.0	4.0	3.1
Total Saline Dominant Species:	0.7	0.8	12.7	8.7	4.5	3.5
<u>Brackish Marsh Vegetation</u>						
Alkali Bulrush	95.4	79.9	84.8	73.3	63.0	76.1
Peppergrass	13.5	26.8	13.6	15.6	7.0	23.6
Spearscale	0.0	0.0	0.0	9.0	6.3	0.0
Total Brackish Dominant Species:	108.9	106.7	98.4	97.9	76.2	99.7
<u>Freshwater Marsh Vegetation</u>						
California Bulrush	0.0	0.0	1.3	4.3	26.7	7.0
Cattail	0.0	0.0	0.1	0.2	1.8	1.1
Total Freshwater Dominant Species:	0.0	0.0	1.4	4.5	28.5	8.1
Total Segment Acreage	109.6	107.5	113.0	115.5	117.5	119.4

Table B12. Acreage Summary of Segment 14 for 1989, 1994/1995, 1996, 1997, 1998 and 1999.
DOMINANT SPECIES CATEGORY

<u>Salt Marsh Vegetation</u>	<u>Year</u>					
	1989	1994/1995	1996	1997	1998	1999
Pickleweed	5.8	8.9	3.4	2.5	0.5	0.8
Cordgrass	2.9	2.0	1.5	2.1	2.0	2.4
Pickleweed-Cordgrass Mix	0.0	0.0	0.0	0.0	0.0	0.0
Pickleweed-Spearscale Mix	0.0	0.0	0.0	0.0	0.0	0.0
Alkali Heath	0.0	0.0	0.0	0.0	0.0	0.0
Gumplant	0.0	0.0	0.0	0.0	0.0	0.0
Peripheral Halophytes	0.7	0.0	0.0	0.6	0.9	1.4
Total Saline Dominant Species:	9.4	10.9	4.9	5.2	3.4	4.6
<u>Brackish Marsh Vegetation</u>						
Alkali Bulrush	10.8	9.1	14.6	16.7	19.3	18.5
Peppergrass	0.0	0.1	0.5	0.3	0.1	0.4
Spearscale	0.0	0.0	0.0	0.0	0.1	0.0

Total Brackish Dominant Species:	10.8	9.2	15.1	17.0	19.4	18.9
<u>Freshwater Marsh Vegetation</u>						
California Bulrush	0.0	0.0	0.0	0.0	0.0	0.0
Cattail	0.0	0.0	0.0	0.0	0.0	0.0
Total Freshwater Dominant Species:	0.0	0.0	0.0	0.0	0.0	0.0
Total Segment Acreage	20.2	20.1	20.0	22.2	23.0	23.9

Table B13. Acreage Summary of Segment 15 for 1989, 1994/1995, 1996, 1997, 1998 and 1999.
DOMINANT SPECIES CATEGORY

	<u>Year</u>					
	1989	1994/1995	1996	1997	1998	1999
<u>Salt Marsh Vegetation</u>						
Pickleweed	1.3	4.2	2.0	1.2	0.4	0.2
Cordgrass	0.8	0.7	0.4	0.7	0.2	0.8
Pickleweed-Cordgrass Mix	0.0	0.0	0.0	0.0	0.0	0.0
Pickleweed-Spearscale Mix	0.0	0.0	0.0	0.0	0.0	0.0
Alkali Heath	0.0	0.0	0.0	0.0	0.0	0.0
Gumplant	0.0	0.0	0.0	0.0	0.0	0.0
Peripheral Halophytes	0.2	0.0	0.2	0.5	0.8	1.4
Total Saline Dominant Species:	2.3	4.9	2.6	2.3	1.3	2.4
<u>Brackish Marsh Vegetation</u>						
Alkali Bulrush	20.1	16.7	18.7	17.9	22.5	21.0
Peppergrass	7.5	7.8	7.4	8.9	6.1	9.8
Spearscale	0.0	0.0	0.0	0.3	0.7	0.2
Total Brackish Dominant Species:	27.6	24.5	26.1	27.2	29.2	31.0
<u>Freshwater Marsh Vegetation</u>						
California Bulrush	0.0	0.0	0.0	0.0	0.0	0.0
Cattail	0.0	0.0	0.0	0.0	0.0	0.0
Total Freshwater Dominant Species:	0.0	0.0	0.0	0.0	0.0	0.0
Total Segment Acreage	29.9	29.4	28.7	29.5	30.5	33.4

Table B14. Acreage Summary of Segment 16 for 1989, 1994/1995, 1996, 1997, 1998 and 1999.
DOMINANT SPECIES CATEGORY

	<u>Year</u>					
	1989	1994/1995	1996	1997	1998	1999
<u>Salt Marsh Vegetation</u>						
Pickleweed	0.0	0.1	0.0	0.0	0.0	0.0
Cordgrass	0.0	0.0	0.0	0.0	0.0	0.0
Pickleweed-Cordgrass Mix	0.0	0.0	0.0	0.0	0.0	0.0
Pickleweed-Spearscale Mix	0.0	0.0	0.0	0.0	0.0	0.0
Alkali Heath	0.0	0.0	0.0	0.2	0.0	0.0
Gumplant	0.0	0.0	0.0	0.0	0.0	0.0
Peripheral Halophytes	1.0	0.0	2.1	1.1	0.0	0.0
Total Saline Dominant Species:	1.0	0.1	2.1	1.3	0.0	0.0
<u>Brackish Marsh Vegetation</u>						
Alkali Bulrush	36.5	29.4	35.3	18.2	33.6	28.2
Peppergrass	8.7	14.8	5.7	4.0	0.9	12.3
Spearscale	0.0	0.0	0.0	18.4	5.7	0.9
Total Brackish Dominant Species:	45.2	44.2	41.0	40.6	40.2	41.4
<u>Freshwater Marsh Vegetation</u>						
California Bulrush	0.0	0.0	0.3	0.7	0.7	3.4
Cattail	0.0	0.0	0.1	0.1	0.0	0.1
Total Freshwater Dominant Species:	0.0	0.0	0.4	0.9	0.7	3.5

Total Segment Acreage	46.2	44.2	45.1	43.3	42.8	54.8
------------------------------	------	------	------	------	------	------

Table B15. Acreage Summary of Segment 17 for 1989, 1994/1995, 1996, 1997, 1998 and 1999.
DOMINANT SPECIES CATEGORY

	<u>Year</u>					
	1989	1994/1995	1996	1997	1998	1999
<u>Salt Marsh Vegetation</u>						
Pickleweed	1.9	1.8	0.0	0.1	0.0	0.0
Cordgrass	0.0	0.0	0.0	0.0	0.0	0.0
Pickleweed-Cordgrass Mix	0.0	0.0	0.0	0.0	0.0	0.0
Pickleweed-Spearscale Mix	0.0	0.0	0.0	0.0	0.0	0.0
Alkali Heath	0.0	0.0	1.8	2.3	0.0	0.1
Gumplant	0.0	0.0	0.0	0.0	0.0	0.0
Peripheral Halophytes	0.7	0.0	0.0	1.1	2.1	1.8
Total Saline Dominant Species:	2.6	1.8	1.8	3.5	2.1	1.9
<u>Brackish Marsh Vegetation</u>						
Alkali Bulrush	90.7	75.9	75.9	44.5	76.3	68.3
Peppergrass	7.8	18.9	18.9	21.1	11.7	28.4
Spearscale	0.0	0.0	0.0	26.6	11.3	0.0
Total Brackish Dominant Species:	98.5	94.8	94.8	92.2	99.3	96.7
<u>Freshwater Marsh Vegetation</u>						
California Bulrush	0.0	0.0	0.0	0.1	0.1	0.0
Cattail	0.0	0.0	0.0	0.5	0.7	0.2
Total Freshwater Dominant Species:	0.0	0.0	0.0	0.5	0.8	0.2
Total Segment Acreage	101.1	96.6	96.6	98.4	104.2	102.8

Table B16. Acreage Summary of Segment 18 for 1989, 1994/1995, 1996, 1997, 1998 and 1999.
DOMINANT SPECIES CATEGORY

	<u>Year</u>					
	1989	1994/1995	1996	1997	1998	1999
<u>Salt Marsh Vegetation</u>						
Pickleweed	1.1	2.1	0.8	1.6	0.6	0.7
Cordgrass	0.0	0.0	0.0	0.2	0.2	0.0
Pickleweed-Cordgrass Mix	0.0	0.0	0.0	0.0	0.0	0.0
Pickleweed-Spearscale Mix	0.0	0.0	0.9	0.1	1.3	0.0
Alkali Heath	0.0	0.3	0.2	0.3	0.1	0.1
Gumplant	0.0	0.0	0.0	0.0	0.0	0.0
Peripheral Halophytes	0.6	0.0	0.6	1.7	1.3	2.1
Total Saline Dominant Species:	1.7	2.4	2.5	3.8	3.5	2.9
<u>Brackish Marsh Vegetation</u>						
Alkali Bulrush	32.7	24.2	24.7	13.4	24.2	22.9
Peppergrass	3.5	8.2	7.2	4.4	2.3	8.3
Spearscale	0.0	0.0	0.0	12.1	3.7	1.3
Total Brackish Dominant Species:	36.2	32.4	31.9	29.8	30.3	32.5
<u>Freshwater Marsh Vegetation</u>						
California Bulrush	0.0	0.0	0.0	0.1	0.1	0.0
Cattail	0.0	0.0	0.0	0.1	0.2	0.1
Total Freshwater Dominant Species:	0.0	0.0	0.0	0.2	0.3	0.1

Total Segment Acreage	37.9	33.8	38.7	36.8	37.4	38.3
------------------------------	------	------	------	------	------	------

Table B17. Acreage Summary of Segment 19 for 1989, 1994/1995, 1996, 1997, 1998 and 1999.
DOMINANT SPECIES CATEGORY

<u>Salt Marsh Vegetation</u>	<u>Year</u>					
	1989	1994/1995	1996	1997	1998	1999
Pickleweed	6.2	11.3	2.6	2.1	30.9	1.0
Cordgrass	0.0	2.0	1.8	0.7	0.1	0.5
Pickleweed-Cordgrass Mix	0.0	0.0	0.0	0.0	0.0	0.2
Pickleweed-Spearscale Mix	0.0	0.0	0.6	0.0	0.2	0.0
Alkali Heath	0.0	0.4	0.2	0.3	0.0	0.1
Gumplant	0.0	0.0	0.0	0.0	0.0	0.0
Peripheral Halophytes	0.5	0.5	1.5	2.8	3.6	3.8
Total Saline Dominant Species:	6.7	14.2	6.7	6.0	34.8	5.6
<u>Brackish Marsh Vegetation</u>						
Alkali Bulrush	31.4	22.1	31.4	24.7	0.8	29.8
Peppergrass	0.5	1.1	1.7	1.2	0.3	2.0
Spearscale	0.0	0.0	0.0	4.2	0.5	0.1
Total Brackish Dominant Species:	31.9	23.2	33.1	30.1	1.7	31.9
<u>Freshwater Marsh Vegetation</u>						
California Bulrush	0.0	0.0	0.0	0.0	0.0	0.0
Cattail	0.0	0.0	0.0	0.2	0.0	0.6
Total Freshwater Dominant Species:	0.0	0.0	0.0	0.2	0.0	0.6
Total Segment Acreage	38.6	37.1	40.8	36.2	36.5	38.4

Table B18. Acreage Summary of Segment 20 for 1989, 1994/1995, 1996, 1997, 1998 and 1999.
DOMINANT SPECIES CATEGORY

<u>Salt Marsh Vegetation</u>	<u>Year</u>					
	1989	1994/1995	1996	1997	1998	1999
Pickleweed	29.6	31.2	18.6	18.2	14.6	14.4
Cordgrass	0.0	6.0	5.0	4.7	2.7	2.6
Pickleweed-Cordgrass Mix	0.0	0.0	0.0	0.0	0.0	0.0
Pickleweed-Spearscale Mix	0.0	0.0	0.0	0.1	0.2	0.0
Alkali Heath	0.0	0.0	0.0	0.1	0.2	0.0
Gumplant	0.0	0.0	0.0	0.0	0.0	0.0
Peripheral Halophytes	0.3	0.0	1.6	1.4	3.3	1.9
Total Saline Dominant Species:	29.9	37.2	25.2	24.5	20.9	18.9
<u>Brackish Marsh Vegetation</u>						
Alkali Bulrush	25.3	17.0	28.9	33.1	36.4	37.9
Peppergrass	2.0	3.3	2.5	3.3	3.3	6.7
Spearscale	0.0	0.0	0.0	0.1	2.1	0.1
Total Brackish Dominant Species:	27.3	20.3	31.4	36.5	41.8	44.7
<u>Freshwater Marsh Vegetation</u>						
California Bulrush	0.0	0.0	0.0	0.0	0.0	0.0
Cattail	0.0	0.0	0.0	0.0	0.0	0.0
Total Freshwater Dominant Species:	0.0	0.0	0.0	0.0	0.0	0.0

Total Segment Acreage	57.2	57.5	56.7	61.1	62.7	63.6
------------------------------	------	------	------	------	------	------

Table B19. Acreage Summary of Segment 21 for 1989, 1994/1995, 1996, 1997, 1998 and 1999.

DOMINANT SPECIES CATEGORY

	<u>Year</u>					
	1989	1994/1995	1996	1997	1998	1999
<u>Salt Marsh Vegetation</u>						
Pickleweed	2.9	7.0	2.9	2.2	1.1	1.0
Cordgrass	0.4	0.4	0.3	0.4	0.3	0.2
Pickleweed-Cordgrass Mix	0.0	0.0	0.0	0.0	0.0	0.0
Pickleweed-Spearscale Mix	0.0	0.0	0.0	0.0	0.0	0.0
Alkali Heath	0.0	0.0	0.0	0.1	0.0	0.0
Gumplant	0.0	0.0	0.0	0.0	0.0	0.0
Peripheral Halophytes	0.8	3.6	0.4	0.3	1.2	0.9
Total Saline Dominant Species:	4.1	11.0	3.6	2.9	2.7	2.1
<u>Brackish Marsh Vegetation</u>						
Alkali Bulrush	20.0	15.1	18.6	17.6	20.6	20.5
Peppergrass	3.0	3.7	4.1	5.3	3.4	6.2
Spearscale	0.0	0.0	0.0	0.8	1.0	0.2
Total Brackish Dominant Species:	23.0	18.8	22.7	23.7	24.9	26.9
<u>Freshwater Marsh Vegetation</u>						
California Bulrush	0.0	0.0	0.0	0.0	0.0	0.0
Cattail	0.0	0.0	0.0	0.0	0.0	0.0
Total Freshwater Dominant Species:	0.0	0.0	0.0	0.0	0.0	0.0
Total Segment Acreage	27.1	29.8	26.7	26.7	27.6	29.0

Table B20. Acreage Summary of Segment 22 for 1989, 1994/1995, 1996, 1997, 1998 and 1999.

DOMINANT SPECIES CATEGORY

	<u>Year</u>					
	1989	1994/1995	1996	1997	1998	1999
<u>Salt Marsh Vegetation</u>						
Pickleweed	5.2	6.1	7.3	6.1	5.2	5.0
Cordgrass	2.3	3.9	2.8	3.8	3.5	4.7
Pickleweed-Cordgrass Mix	0.0	0.0	0.0	0.0	0.0	0.1
Pickleweed-Spearscale Mix	0.0	0.0	0.0	0.0	0.0	0.0
Alkali Heath	0.0	0.0	0.0	0.0	0.0	0.0
Gumplant	0.0	0.0	0.0	0.0	0.0	0.0
Peripheral Halophytes	0.4	0.0	0.5	1.0	1.2	0.9
Total Saline Dominant Species:	7.9	10.0	10.6	10.9	9.9	10.7
<u>Brackish Marsh Vegetation</u>						
Alkali Bulrush	0.0	0.0	0.2	1.0	2.9	2.7
Peppergrass	0.0	0.2	0.4	0.0	0.0	0.6
Spearscale	0.0	0.0	0.0	0.0	0.0	0.0
Total Brackish Dominant Species:	0.0	0.2	0.6	1.0	2.9	3.3
<u>Freshwater Marsh Vegetation</u>						
California Bulrush	0.0	0.0	0.0	0.0	0.0	0.0
Cattail	0.0	0.0	0.0	0.0	0.0	0.0

Total Freshwater Dominant Species:	0.0	0.0	0.0	0.0	0.0	0.0
Total Segment Acreage	7.9	10.2	11.2	11.9	12.8	14.1

Table B21. Acreage Summary of Segment 23 for 1989, 1994/1995, 1996, 1997, 1998 and 1999.
DOMINANT SPECIES CATEGORY

	<u>Year</u>					
	1989	1994/1995	1996	1997	1998	1999
<u>Salt Marsh Vegetation</u>						
Pickleweed	8.9	14.1	14.1	11.1	10.2	10.2
Cordgrass	8.5	3.7	3.6	4.8	6.2	5.9
Pickleweed-Cordgrass Mix	0.0	0.0	0.0	0.1	0.0	1.3
Pickleweed-Spearscale Mix	0.0	0.0	0.0	0.0	0.0	0.0
Alkali Heath	0.0	0.0	0.2	0.0	0.0	0.2
Gumplant	0.0	0.0	0.0	0.0	0.0	0.0
Peripheral Halophytes	0.3	0.0	0.8	1.4	1.7	1.5
Total Saline Dominant Species:	17.7	17.8	18.7	17.4	18.1	19.1
<u>Brackish Marsh Vegetation</u>						
Alkali Bulrush	0.0	0.0	0.0	0.0	0.0	0.0
Peppergrass	0.0	0.0	0.0	0.0	0.0	0.0
Spearscale	0.0	0.0	0.0	0.0	0.0	0.0
Total Brackish Dominant Species:	0.0	0.0	0.0	0.0	0.0	0.0
<u>Freshwater Marsh Vegetation</u>						
California Bulrush	0.0	0.0	0.0	0.0	0.0	0.0
Cattail	0.0	0.0	0.0	0.0	0.0	0.0
Total Freshwater Dominant Species:	0.0	0.0	0.0	0.0	0.0	0.0
Total Segment Acreage	17.7	17.8	21.2	17.7	18.4	19.8

Table B22. Acreage Summary of Segment 24* for 1994/1995, 1996, 1997, 1998 and 1999.
DOMINANT SPECIES CATEGORY

	<u>Year</u>				
	1994/1995	1996	1997	1998	1999
<u>Salt Marsh Vegetation</u>					
Pickleweed	0.8	0.2	0.6	0.6	0.2
Cordgrass	0.0	0.0	0.0	0.0	0.0
Pickleweed-Cordgrass Mix	0.0	0.0	0.0	0.0	0.0
Pickleweed-Spearscale Mix	0.0	0.0	0.0	0.0	0.0
Alkali Heath	0.0	0.0	0.0	0.0	0.0
Gumplant	0.0	0.0	0.0	0.0	0.0
Peripheral Halophytes	1.5	2.2	0.7	0.8	0.5
Total Saline Dominant Species:	2.3	2.4	1.3	1.4	0.7
<u>Brackish Marsh Vegetation</u>					
Alkali Bulrush	1.5	2.0	1.8	2.2	2.4
Peppergrass	7.0	6.0	5.7	7.1	7.1
Spearscale	0.0	0.0	0.0	0.5	0.1
Total Brackish Dominant Species:	8.5	8.0	7.5	9.7	9.6
<u>Freshwater Marsh Vegetation</u>					
California Bulrush	1.4	1.6	1.9	2.0	2.6

Cattail	0.0	0.0	0.0	0.0	0.0
Total Freshwater Dominant Species:	1.4	1.6	1.9	2.0	2.6
Total Segment Acreage	20.2	12.1	10.7	11.2	12.9

* Segment 24 not mapped in 1989

Table B23. Acreage Summary of Segment 25* for 1994/1995, 1996, 1997, 1998 and 1999.

DOMINANT SPECIES CATEGORY

	Year				
	1994/1995	1996	1997	1998	1999
<u>Salt Marsh Vegetation</u>					
Pickleweed	0.0	0.0	0.0	0.0	0.0
Cordgrass	0.0	0.0	0.0	0.0	0.0
Pickleweed-Cordgrass Mix	0.0	0.0	0.0	0.0	0.0
Pickleweed-Spearscale Mix	0.0	0.0	0.0	0.0	0.0
Alkali Heath	0.0	0.0	0.0	0.0	0.0
Gumplant	0.0	0.0	0.0	0.0	0.0
Peripheral Halophytes	5.3	4.0	2.6	0.0	1.0
Total Saline Dominant Species:	5.3	4.0	2.6	0.0	1.0
<u>Brackish Marsh Vegetation</u>					
Alkali Bulrush	2.9	4.3	3.4	3.3	5.8
Peppergrass	10.4	7.7	6.5	48.6	7.6
Spearscale	0.0	0.0	0.3	0.5	0.1
Total Brackish Dominant Species:	13.3	12.0	10.3	52.3	13.5
<u>Freshwater Marsh Vegetation</u>					
California Bulrush	29.8	30.3	31.3	0.1	38.6
Cattail	0.2	0.8	1.5	0.2	2.0
Total Freshwater Dominant Species:	30.0	31.1	32.8	0.3	40.6
Total Segment Acreage	48.6	47.2	46.2	51.7	55.3

*Segment 25 not mapped in 1989

Table B24. Acreage Summary of Segment 26* for 1994/1995, 1996, 1997, 1998 and 1999.

DOMINANT SPECIES CATEGORY

	Year				
	1994/1995	1996	1997	1998	1999
<u>Salt Marsh Vegetation</u>					
Pickleweed	0.0	0.0	0.0	0.0	0.0
Cordgrass	0.0	0.0	0.0	0.0	0.0
Pickleweed-Cordgrass Mix	0.0	0.0	0.0	0.0	0.0
Pickleweed-Spearscale Mix	0.0	0.0	0.0	0.0	0.0
Alkali Heath	0.0	0.0	0.0	0.0	0.0
Gumplant	0.0	0.0	0.0	0.0	0.0
Peripheral Halophytes	1.3	1.3	0.8	0.1	0.0
Total Saline Dominant Species:	1.3	1.3	0.8	0.1	0.0
<u>Brackish Marsh Vegetation</u>					
Alkali Bulrush	0.0	0.0	0.0	0.0	0.0
Peppergrass	2.5	2.6	0.6	0.1	2.9
Spearscale	0.0	0.0	0.0	0.0	0.0
Total Brackish Dominant Species:	2.5	2.6	0.6	0.2	3.0
<u>Freshwater Marsh Vegetation</u>					

California Bulrush	17.8	18.7	17.5	18.8	18.0
Cattail	0.1	0.2	0.4	0.3	0.1
Total Freshwater Dominant Species:	17.9	18.9	17.9	19.1	18.1
Total Segment Acreage	23.7	23.1	23.0	24.1	23.7

*Segment 26 not mapped in 1989

Table B25. Acreage Summary of Segment 27* for 1996, 1997, 1998 and 1999.

DOMINANT SPECIES CATEGORY

	<u>Year</u>			
	1996	1997	1998	1999
<u>Salt Marsh Vegetation</u>				
Pickleweed	0.0	0.9	0.0	0.0
Cordgrass	0.0	0.0	0.0	0.0
Peripheral Halophytes	1.0	2.1	2.3	0.0
Total Saline Dominant Species:	1.0	3.0	2.3	0.0
<u>Brackish Marsh Vegetation</u>				
Alkali Bulrush	11.4	9.1	8.9	7.4
Peppergrass	0.6	1.7	0.1	1.2
Spearscale**	0.0	0.1	0.1	0.0
Total Brackish Dominant Species:	12.0	10.8	9.1	8.6
<u>Freshwater Marsh Vegetation</u>				
California Bulrush	3.3	4.4	6.7	4.7
Cattail	7.6	7.8	8.4	10.8
Total Freshwater Dominant Species:	10.9	12.2	15.2	15.5
Total Segment Acreage	35.0	35.7	35.7	36.5

*Segment 27 not mapped in 1989 and

1994/1995

**Not a Dominant Species Category in 1996

Table B26. Acreage Summary of Segment 28* for 1989, 1996, 1997, 1998 and 1999.

DOMINANT SPECIES CATEGORY

	<u>Year</u>				
	1989	1996	1997	1998	1999
<u>Salt Marsh Vegetation</u>					
Pickleweed	0.0	0.5	0.2	0.1	0.1
Cordgrass	8.9	1.6	1.8	0.8	0.0
Peripheral Halophytes	0.0	0.3	1.4	4.0	3.4
Total Saline Dominant Species:	8.9	2.4	3.4	4.8	3.5
<u>Brackish Marsh Vegetation</u>					
Alkali Bulrush	48.7	53.7	49.8	61.9	57.0
Peppergrass	7.4	9.9	15.8	2.2	10.2
Spearscale**	0.0	0.0	0.1	0.2	0.0
Total Brackish Dominant Species:	56.1	63.5	65.7	64.3	67.2
<u>Freshwater Marsh Vegetation</u>					
California Bulrush	0.2	10.5	9.1	15.5	15.6
Cattail	0.0	0.3	0.4	0.5	0.6
Total Freshwater Dominant Species:	0.2	10.8	9.5	16.0	16.2
Total Segment Acreage	65.2	77.8	78.9	85.7	90.3

*Segment 28 not mapped in 1994/1995

**Not a Dominant Species Category in 1996

**Table B27. Acreage Summary of Segment 29* for 1989, 1996, 1997, 1998 and 1999.
DOMINANT SPECIES CATEGORY**

	<u>Year</u>				
	1989	1996	1997	1998	1999
<u>Salt Marsh Vegetation</u>					
Pickleweed	21.9	14.8	12.1	9.0	9.3
Cordgrass	12.9	5.6	6.8	4.6	2.3
Peripheral Halophytes	0.0	2.2	4.3	5.8	5.6
Total Saline Dominant Species:	34.8	22.5	23.2	19.4	17.2
<u>Brackish Marsh Vegetation</u>					
Alkali Bulrush	18.4	48.4	47.2	58.7	65.5
Peppergrass	9.0	10.0	9.5	3.9	11.0
Spearscale**	0.0	0.0	0.3	0.1	0.1
Total Brackish Dominant Species:	27.4	58.3	57.0	62.6	76.6
<u>Freshwater Marsh Vegetation</u>					
California Bulrush	0.0	0.0	0.0	0.0	0.3
Cattail	0.0	0.0	0.0	0.0	0.0
Total Freshwater Dominant Species:	0.0	0.0	0.0	0.0	0.3
Total Segment Acreage	72.2	81.1	80.6	82.0	94.3

*Segment 29 not mapped in 1994/1995

**Not a Dominant Species Category in 1996

**Table B28. Acreage Summary of Segment 30* for 1989, 1996, 1997, 1998 and 1999.
DOMINANT SPECIES CATEGORY**

	<u>Year</u>				
	1989	1996	1997	1998	1999
<u>Salt Marsh Vegetation</u>					
Pickleweed	24.9	26.5	23.1	19.7	21.0
Cordgrass	6.1	8.0	9.8	10.7	13.0
Peripheral Halophytes	2.6	1.5	2.6	2.9	3.7
Total Saline Dominant Species:	33.6	36.0	35.5	33.3	37.7
<u>Brackish Marsh Vegetation</u>					
Alkali Bulrush	0.0	1.5	1.7	6.5	5.5
Peppergrass	1.2	2.0	0.0	0.0	0.0
Spearscale**	0.0	0.0	0.0	0.0	0.0
Total Brackish Dominant Species:	1.2	3.4	1.7	6.5	5.5
<u>Freshwater Marsh Vegetation</u>					
California Bulrush	0.0	0.0	0.0	0.0	0.0
Cattail	0.0	0.0	0.0	0.0	0.0
Total Freshwater Dominant Species:	0.0	0.0	0.0	0.0	0.0

Total Segment Acreage	34.8	39.4	37.7	40.8	46.8
------------------------------	------	------	------	------	------

***Segment 30 not mapped in 1994/1995**

****Not a Dominant Species Category in 1996**

**APPENDIX C
SOUTH BAY MARSHES:
PLANT LIST**

Appendix G. Plants Observed in the South Bay Marsh Project Site		
FAMILY NAME	SCIENTIFIC NAME	COMMON NAME
Aceraceae	<i>Acer negundo</i> ssp. <i>californica</i>	California box elder
Aizoaceae	<i>Mesembryanthemum nodiflorum</i>	slender-leaved iceplant
	<i>Tetragonia tetragonioides</i>	New Zealand spinach
Apiaceae	<i>Foeniculum vulgare</i>	sweet fennel
	<i>Conium maculatum</i>	poison hemlock
Asteraceae	<i>Baccharis pilularis</i>	coyote brush
	<i>Carduus pycnocephalus</i>	Italian thistle
	<i>Centaurea solstitialis</i>	yellow star-thistle
	<i>Conyza canadensis</i>	horsetail
	<i>Grindelia</i> sp.	gumplant
	<i>Picris echioides</i>	bristly ox-tongue
Brassicaceae	<i>Brassica nigra</i>	black mustard
	<i>Hirschfeldia incana</i>	small-pod mustard
	<i>Lepidium latifolium</i>	perennial peppergrass
Chenopodiaceae	<i>Atriplex semibaccata</i>	Australian saltbush
	<i>Atriplex triangularis</i>	spearscale
	<i>Bassia hyssopifolia</i>	five-hook bassia
	<i>Salicornia virginica</i>	common pickleweed
	<i>Salicornia europaea</i>	annual pickleweed
Cuscutaceae	<i>Cuscuta salina</i> var. <i>major</i>	salt marsh dodder
Cyperaceae	<i>Scirpus acutus</i>	tule
	<i>Scirpus californicus</i>	California bulrush
	<i>Scirpus maritimus</i>	alkali bulrush
Frankeniaceae	<i>Frankenia salina</i>	alkali heath
Juglandaceae	<i>Juglans californica</i>	California black walnut
Poaceae	<i>Bromus diandrus</i>	ripgut grass
	<i>Bromus hordeaceus</i>	soft chess
	<i>Distichlis spicata</i>	saltgrass
	<i>Hordeum</i> sp.	barley
	<i>Spartina foliosa</i>	cordgrass
Polygonaceae	<i>Polygonum punctatum</i>	knotweed
Salicaceae	<i>Populus fremontii</i>	Fremont's cottonwood
Solanaceae	<i>Solanum americanum</i>	deadly nightshade
	<i>Nicotiana glauca</i>	tree-tobacco
Typhaceae	<i>Typha</i> sp.	cattail

The species are arranged alphabetically by family name for all vascular plants encountered during the plant survey. Plants are also listed alphabetically within each family. In some cases it was not possible to accurately identify a particular plant to the species level due to the absence of specific anatomic structures required for identification.

